

## INTERNATIONAL COOPERATION TREATY

PCT

## NOTIFICATION OF ELECTION

(PCT Rule 61.2)

From the INTERNATIONAL BUREAU

To:

Assistant Commissioner for Patents  
 United States Patent and Trademark  
 Office  
 Box PCT  
 Washington, D.C.20231  
 ÉTATS-UNIS D'AMÉRIQUE

in its capacity as elected Office

<b>Date of mailing (day/month/year)</b> 14 September 1999 (14.09.99)	
<b>International application No.</b> PCT/EP98/08319	<b>Applicant's or agent's file reference</b> B 3121 PCT
<b>International filing date (day/month/year)</b> 18 December 1998 (18.12.98)	<b>Priority date (day/month/year)</b> 23 January 1998 (23.01.98)
<b>Applicant</b> FLEGEL, Willy, A. et al	

1. The designated Office is hereby notified of its election made:

☒ in the demand filed with the International Preliminary Examining Authority on:

02 August 1999 (02.08.99)

☐ in a notice effecting later election filed with the International Bureau on:

2. The election ☒ was  
☐ was not

made before the expiration of 19 months from the priority date or, where Rule 32 applies, within the time limit under Rule 32.2(b).

The International Bureau of WIPO 34, chemin des Colombettes 1211 Geneva 20, Switzerland  Facsimile No.: (41-22) 740.14.35	Authorized officer  F. Baechler  Telephone No.: (41-22) 338.83.38
---	---

## CLAIMS

1. A nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, in its transmembrane and/or intracellular regions.
2. A nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule
- (a) carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, in amino acid positions 2-16, 114-149, 179-225 or/and 267 to 397 with the proviso that said D antigen does carry not a single missense mutation leading to a substitution of phenylalanine in amino acid position 223 by valine or of threonine in position 283 by isoleucine; or
  - (b) carrying a gene conversion involving exons 6 to 9 which are replaced by the corresponding exons of the *RHCE* gene.
3. The nucleic acid molecule of claim 1 or 2 wherein said missense mutation causes an amino acid substitution in position 3, 10, 16, 114, 149, 182, 198, 201, 220, 223, 270, 276, 277, 282, 294, 295, 307, 339, 385 or 393 or a combination of/or involving said substitutions.
4. The nucleic acid molecule of claim 3 wherein said amino acid substitution in position 3 is from Ser to Cys, in position 10 from Arg to Gln, in position 16 from Trp to Cys, in position 114 from Arg to Trp, in position 149 from Ala to Asp, in position 182 from Ser to Thr, in position 198 from Lys to Asn, in position 201 from Thr to Arg, in position 220 from Trp to Arg, in position 223 from Phe to Val, in position 270 from Val to Gly, in position 276 from Ala to Pro, in position 277 from Gly to Glu, in position 282 from Gly to Asp, in position 294 from Ala to Pro, in position 295 from Met to Ile, in position 307 from Gly to Arg, in

position 339 from Gly to Glu, in position 385 from Gly to Ala and in position 393 from Trp to Arg.

5. The nucleic acid molecule of any one of claims 1 to 4 wherein said missense mutation occurs in nucleotide position 8, 29, 48, 340, 446, 544, 594, 602, 658, 667, 809, 819, 826, 830, 845, 880, 885, 919, 1016, 1154 or 1177 or in a combination of said positions.
6. The nucleic acid molecule of claim 5 wherein said missense mutation in position 8 is from C to G, in position 29 from G to A, in position 48 from G to C, in position 340 from C to T, in position 446 from C to A, in position 544 from T to A, in position 594 from A to T, in position 602 from C to G, in position 658 from T to C, in position 667 from T to G, in position 809 from T to G, in position 819 from G to A, in position 826 from G to C, in position 830 from G to A, in position 845 from G to A, in position 880 from G to C, in position 885 from G to T, in position 919 from G to A, in position 1016 from G to A, in position 1154 from G to C and in position 1177 from T to C.
7. The nucleic acid molecule of claim 3 or 4 wherein said combination of substitutions is in positions 182, 198 and 201 and is preferably S182T, K198N, T201R or in position 201 and 223 and is preferably T201R and F223V, or in position 16, 201 and 223 and is preferably W16C, T201R and F223V.
8. The nucleic acid molecule of claim 5 or 6 wherein said combination of missense mutations comprises positions 544, 594 and 602 and is preferably TTTA at position 544, ATTT at position 594 and CTTG at position 602 or comprises positions 602, 677 and 819 and is preferably CTTG at position 602, TTTG at position 667 and GTTA at position 819 or comprises positions 48, 602, 667 and 819 and is preferably GTTC at position 48, CTTG at position 602, TTTG at position 667 and GTTA at position 819.
9. The nucleic acid molecule of any one of claims 1 to 8 which is mRNA or genomic DNA.

ART 34 AMDT

10. A vector comprising the nucleic acid molecule of any one of claims 1 to 9.
11. A non-human host transformed with the vector of claim 10.
12. A method of producing a Rhesus D antigen contributing to the weak D phenotype comprising culturing the host of claim 11 under suitable conditions and isolating the Rhesus D antigen produced.
13. A Rhesus D antigen encoded by the nucleic acid molecule of any one of claims 1 to 9 or produced by the method of claim 12.
14. An oligonucleotide hybridizing under stringent conditions to a portion of the nucleic acid molecule of any one of claims 1 to 9 comprising said at least one missense mutation or to the complementary portion thereof or hybridizing to a region involving the breakpoint of the gene conversion identified in claim 2.
15. An antibody or aptamer or phage specifically binding to the Rhesus D antigen of claim 13.
16. An antibody or aptamer or phage specifically binding to the wild-type Rhesus D antigen or to aberrant Rhesus D antigens but not to the Rhesus D antigen of claim 13.
17. A method for testing for the presence of a nucleic acid molecule encoding a Rhesus D antigen contributing to the weak D phenotype in a sample comprising hybridizing the oligonucleotide of claim 14 under stringent conditions to nucleic acid molecules comprised in the sample obtained from human and detecting said hybridization.
18. The method of claim 17 further comprising digesting the product of said hybridization with a restriction endonuclease and analyzing the product of said digestion.

sub  
C4

19. A method of testing for the presence of a nucleic acid encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising determining the nucleic acid sequence of at least a portion of the nucleic acid molecule of any one of claims 1 to 9, said portion encoding at least one of said missense mutations or a region involving the breakpoint of said gene conversion or of a nucleic acid molecule encoding a Rhesus D antigen contributing to the weak D phenotype, said nucleic acid molecule carrying a combination of missense mutations, as compared to the wild type Rhesus D antigen, one of said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T.
20. The method of claim 19 further comprising, prior to determining said nucleic acid sequence, amplification of at least said portion of said nucleic acid molecule.
21. A method for testing for the presence of a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising carrying out an amplification reaction wherein at least one of the primers employed in said amplification reaction is the oligonucleotide of claim 14, comprising assaying for an amplification product.
22. The method of claim 20 or 21 wherein said amplification is effected by or said amplification is the polymerase chain reaction (PCR).
23. A method for testing for the presence of a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising assaying a sample obtained from a human for specific binding to the antibody or aptamer or phage of claim 15.

24. A method of testing a sample for the presence of wild type Rhesus D antigen and the absence of the Rhesus D antigen of claim 13 comprising assaying a sample obtained from a human for specific binding to an antibody or aptamer or phage of claim 16.
25. The method of any one of claims 17 to 24 wherein said sample is blood, serum, plasma, fetal tissue, saliva, urine, mucosal tissue, mucus, vaginal tissue, fetal tissue obtained from the vagina, skin, hair, hair follicle or another human tissue.
26. The method of 25 comprising enrichment of fetal cells or extraction of fetal DNA or mRNA from material tissue, like peripheral blood, serum or plasma.
27. The method of any one of claims 17 to 26 wherein said nucleic acid molecule or proteinaceous material from said sample is fixed to a solid support.
28. The method of claim 27 wherein said solid support is a chip.
29. Use of the nucleic acid molecule of any one of claims 1 to 9 or of a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying a combination of missense mutations, as compared to the wild type Rhesus D antigen, one of said missense mutations causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T or of a combination thereof for the analysis of a weak Rhesus D phenotype.
30. Use of the nucleic acid molecule of any one of claims 1 to 9, the vector of claim 10 or the Rhesus D antigen of claim 13 for the assessment of the affinity, avidity and/or reactivity of monoclonal antibodies or of polyclonal antisera preferably anti-D antisera, anti-globulin or anti-human-globulin antisera.

31. A method for the characterization of the monoclonal antibodies or polyclonal antisera or of a preparation thereof said method comprising
- (a) testing the nucleic acid of sample of a proband for the presence of a mutation as defined in any one of claims 1 to 8;
  - (b) correlating, on the basis of the mutation status and the allelic status of the RHD gene, the nucleic acid with the RhD antigen density on the surface of red blood cells of said proband;
  - (c) reacting said monoclonal antibodies or polyclonal antisera or said preparation thereof with a cell carrying the RhD antigen on its surface;
  - (d) characterizing said monoclonal antibodies or polyclonal antisera or said preparation thereof on the basis of the results obtained in step (c).
32. The method of claim 31 wherein said characterization comprises the determination of reactivity, sensitivity, avidity, affinity, specificity and/or other characteristics of antibodies and antisera.
33. The method of claim 31 or 32 wherein said cell carrying the RhD antigen on its surface is a red blood cell.
34. Use of an aptamer, phage, monoclonal antibody or a polyclonal antisera as characterized in claims 15 or 16 or a preparation thereof for RhD antigen determination.
35. Use of claim 34 wherein said RhD antigen determination is effected in connection with blood group typing.
36. A preparation comprising the antibody or aptamer or phage of claims 15 or 16.
37. A method of identifying an antibody  $V_H$  or  $V_L$  chain or a combination thereof or an aptamer specifically binding to a weak D antigen of claim 13 comprising
- (a) contacting the weak D antigen of claim 13 with a phage library displaying  $V_H$  or  $V_L$  chains or combinations thereof on the surface of the phage or with aptamers;

- (b) identifying phage or aptamers that bind to said weak D antigen; and optionally
  - (c) repeating steps (a) and (b) one or more times.
38. A method of identifying a monoclonal antibody specifically binding to a weak D polypeptide/antigen of claim 13 comprising
- (a) contacting the weak D polypeptide of claim 13 with one or more monoclonal antibodies;
  - (b) identifying monoclonal antibodies that bind to said weak D polypeptide; and optionally
  - (c) repeating steps (a) and (b) one or more times.
39. A method of identifying an antibody V<sub>H</sub> or V<sub>L</sub> chain or a combination thereof or an aptamer specifically binding to a weak D polypeptide/antigen of claim 13 comprising
- (a) contacting the weak D polypeptide and
    - (aa) a second or more weak D polypeptide(s) and/or
    - (ab) a normal D polypeptidewherein the second or more weak D polypeptide(s) and/or the normal D polypeptide are present in a molar mass that is higher, equal or less than the weak D polypeptide of (a) with a phage library displaying V<sub>H</sub> or V<sub>L</sub> chains or combinations thereof on the surface of the phage or with aptamers;
  - (b) identifying phage or aptamers that bind to said weak D polypeptide of (a); and optionally
  - (c) repeating steps (a) and (b) one or more times.
40. A method of identifying a monoclonal antibody specifically binding to a weak D polypeptide/antigen of claim 13 comprising
- (a) contacting the weak D polypeptide and
    - (aa) a second or more weak D polypeptide(s) and/or.
    - (ab) a normal D polypeptidewherein the second or more weak D polypeptide(s) and/or the normal D polypeptide are present in a molar mass which is higher, equal or less





- 5th  
CS

- Figure 1**

add  
c6

ART 34 AMDT

(1996), 714-716; WO94/29469; WO 97/00957 or Schaper, Current Opinion in Biotechnology 7 (1996), 635-640, and references cited therein. The polynucleotides and vectors of the invention may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Preferably, said cell is a germ line cell, embryonic cell, or egg cell or derived therefrom, most preferably said cell is a stem cell.

Additionally, the invention relates to a host transformed with the vector of the invention.

Appropriate hosts comprise transgenic non human animals, cells such as bacteria, yeast cells, animal, preferably mammalian, fungal cells or insect cells. Transformation protocols including transfection, microinjection, electroporation, etc., are also well known in the art.

Further, the invention relates to a method of producing a Rhesus D antigen contributing to the weak D phenotype comprising culturing the host of the invention under suitable conditions and isolating the Rhesus D antigen produced.

It is preferred that the antigen is exported into the culture medium where it can be collected according to conventions/methods. The term "culturing" as used in accordance with the present invention also comprises the raising of transgenic animals. Using appropriate vectors constructions and optionally appropriate feeds, the antigen may, e.g., be isolated from milk of, e.g. transgenic cows.

The invention additionally relates to Rhesus D antigen encoded by the nucleic acid molecule of the invention or produced by the method of the invention.

Preferably, the antigen is in the same way post transitionally modified and has the same chemical structure as naturally occurring antigen. Accordingly, said antigen, when produced by the method of the invention, is preferably produced in human cells.

Furthermore, the invention relates to an oligonucleotide hybridizing under stringent conditions to a portion of the nucleic acid molecule of the invention comprising said at least one missense mutation or to the complementary portion

be characterized on the basis of the teachings of the present invention. An appropriately characterized anti-D monoclonal antibody can be conveniently used in RhD diagnostics. For example, a suitably characterized monoclonal antibody will be useful in determining the weak D antigen density on the surface of blood cells. Cut-off values for monoclonal antibodies useful in diagnosis can thus be established. This is important for the quality control of antibodies used in RhD diagnosis.

Thus, the invention also relates to a method for the characterization of monoclonal antibodies or polyclonal antisera or of a preparation thereof, said method comprising

- (a) testing the nucleic acid of sample of a proband for the presence of a mutation as defined in accordance with the invention;
- (b) correlating, on the basis of the mutation status and the allelic status of the *RHD* gene, the nucleic acid with the RhD antigen density on the surface of red blood cells of said proband;
- (c) reacting said monoclonal antibodies or polyclonal antisera or said preparation thereof with a cell carrying the RhD antigen on its surface;
- (d) characterizing said monoclonal antibodies or polyclonal antisera or said preparation thereof on the basis of the results obtained in step (c).

As regards the term "allelic status", this term describes the possibilities that the *RHD* alleles in a proband are present in a homozygous, heterozygous or hemizygous state. Also comprised by this term is the possibility that the two alleles carry two different mutations (including the conversion) defined herein above.

In a preferred embodiment of the method of the invention, said characterization comprises the determination of reactivity, sensitivity, avidity, affinity, specificity and/or other characteristics of antibodies and antisera.

Furthermore preferred is a method wherein said cell carrying the RhD antigen on its surface is a red blood cell.

# PATENT COOPERATION TREATY

## PCT

REC'D 16 MAY 2000

WIPO

PCT

### INTERNATIONAL PRELIMINARY EXAMINATION REPORT

(PCT Article 36 and Rule 70)

Applicant's or agent's file reference <b>B 3121 PCT</b>	See Notification of Transmittal of International Preliminary Examination Report (Form PCT/IPEA/416) <b>FOR FURTHER ACTION</b>	
International application No. <b>PCT/EP98/08319</b>	International filing date (day/month/year) <b>18/12/1998</b>	Priority date (day/month/year) <b>23/01/1998</b>
International Patent Classification (IPC) or national classification and IPC <b>C12N15/12</b>		
Applicant <b>DRK BLUTSPENDEDIENST BADEN-WÜRTTEMBERG ...et al.</b>		

1. This international preliminary examination report has been prepared by this International Preliminary Examining Authority and is transmitted to the applicant according to Article 36.

2. This REPORT consists of a total of 5 sheets, including this cover sheet.

☒ This report is also accompanied by ANNEXES, i.e. sheets of the description, claims and/or drawings which have been amended and are the basis for this report and/or sheets containing rectifications made before this Authority (see Rule 70.16 and Section 607 of the Administrative Instructions under the PCT).

These annexes consist of a total of 11 sheets.

3. This report contains indications relating to the following items:

- I ☒ Basis of the report
- II ☒ Priority
- III ☐ Non-establishment of opinion with regard to novelty, inventive step and industrial applicability
- IV ☐ Lack of unity of invention
- V ☒ Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement
- VI ☐ Certain documents cited
- VII ☐ Certain defects in the international application
- VIII ☒ Certain observations on the international application

Date of submission of the demand

02/08/1999

Date of completion of this report

11.05.00

Name and mailing address of the international preliminary examining authority:



European Patent Office  
D-80298 Munich  
Tel. +49 89 2399 - 0 Tx: 523656 epmu d  
Fax: +49 89 2399 - 4465

Authorized officer

van Heusden, M

Telephone No. +49 89 2399 8145



**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**

International application No. PCT/EP98/08319

**I. Basis of the report**

1. This report has been drawn on the basis of (*substitute sheets which have been furnished to the receiving Office in response to an invitation under Article 14 are referred to in this report as "originally filed" and are not annexed to the report since they do not contain amendments.*):

**Description, pages:**

1-10,12-18,                      as originally filed  
20-47

11,19                      as received on                      17/04/2000    with letter of                      17/04/2000

**Claims, No.:**

1-48                      as received on                      17/04/2000    with letter of                      17/04/2000

**Drawings, sheets:**

1/5-5/5                      as originally filed

2. The amendments have resulted in the cancellation of:

- ☐ the description,                      pages:  
☐ the claims,                      Nos.:  
☐ the drawings,                      sheets:

3. ☐ This report has been established as if (some of) the amendments had not been made, since they have been considered to go beyond the disclosure as filed (Rule 70.2(c)):

4. Additional observations, if necessary:

**II. Priority**

1. ☐ This report has been established as if no priority had been claimed due to the failure to furnish within the prescribed time limit the requested:
- ☐ copy of the earlier application whose priority has been claimed.
- ☐ translation of the earlier application whose priority has been claimed.

**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT**

International application No. PCT/EP98/08319

2. ☐ This report has been established as if no priority had been claimed due to the fact that the priority claim has been found invalid.

Thus for the purposes of this report, the international filing date indicated above is considered to be the relevant date.

3. Additional observations, if necessary:

**see separate sheet**

**V. Reasoned statement under Article 35(2) with regard to novelty, inventive step or industrial applicability; citations and explanations supporting such statement**

1. Statement

Novelty (N)	Yes:	Claims	1-48
	No:	Claims	
Inventive step (IS)	Yes:	Claims	1-48
	No:	Claims	
Industrial applicability (IA)	Yes:	Claims	1-48
	No:	Claims	

2. Citations and explanations

**see separate sheet**

**VIII. Certain observations on the international application**

The following observations on the clarity of the claims, description, and drawings or on the question whether the claims are fully supported by the description, are made:

**see separate sheet**

**Additional remarks to section II:**

1. The documents mentioned in this IPER are numbered as in the International Search Report (ISR), i.e. D1 corresponds to the first document of the ISR etc.
2. The priority document pertaining to the present application was not available at the time of establishing this IPER. Hence, the current assessment is based on the assumption that all claims enjoy priority rights from the filing date of the priority document. If it later turns out that this is not correct, the document by Legler et al. cited as D4 in the ISR could become relevant to assess whether the subject matter of the present application satisfies the criteria set forth in Article 33(1) PCT.

**Additional remarks to section V:**

**1. Novelty and Inventive step (Article 33(2) and (3) PCT)**

- 1.1 The present application discloses Rhesus D antigens having missense mutations or a gene conversion involving exons 6-9, both resulting in a weak D phenotype, the encoding nucleic acid molecules, antibodies binding said mutated antigens, and methods of testing for the presence of said mutated RhD antigens.
- 1.2 The present application shows for the first time a molecular basis for the weak D-phenotype. None of the cited prior art discloses any molecular basis for the weak D-phenotype, let alone the contribution of at least one missense mutation in the transmembrane and/or intracellular region of the wildtype Rhesus D antigen. In fact D1 states that the weak D-phenotype is not the result of rearrangement of mutation in the coding sequence of the RHD gene and thus teaches away from the present invention. Therefore the subject matter of the present application is considered novel and inventive over the cited prior art and thus complies with the criterion set forth in Article 33(2) and (3) PCT.

**2. Industrial applicability (Article 33(4) PCT)**

The subject matter of claims 1-48 is industrially applicable.



**INTERNATIONAL PRELIMINARY  
EXAMINATION REPORT - SEPARATE SHEET**

---

International application No. PCT/EP98/08319

**Additional remarks to section VIII:**

The following objections are raised under **Article 6 PCT** concerning the clarity of the claims:

1. The wording '**or involving**' in claim 3 is meaningless.
2. The subject matter of claim 14 lacks clarity in that it is not clear whether the word '**comprising**' refers to 'the nucleic acid molecule' or to the 'portion of..' It seems that it is crucial that the oligonucleotide should hybridize to a portion (of the nucleic acid molecule of any of claims 1-9) said portion comprising said at least one missense mutation, which is not clear from the claim as presently worded.

# TENT COOPERATION TREATY

# PCT

## INTERNATIONAL SEARCH REPORT

(PCT Article 18 and Rules 43 and 44)

Applicant's or agent's file reference <b>B3121PCT</b>	<b>FOR FURTHER ACTION</b> <small>see Notification of Transmittal of International Search Report (Form PCT/ISA/220) as well as, where applicable, item 5 below.</small>	
International application No. <b>PCT/EP 98/ 08319</b>	International filing date (day/month/year) <b>18/12/1998</b>	(Earliest) Priority Date (day/month/year) <b>23/01/1998</b>
Applicant  <b>DRK BLUTSPENDEDIENST BADEN-WÜRTTEMBERG ...et al.</b>		

This International Search Report has been prepared by this International Searching Authority and is transmitted to the applicant according to Article 18. A copy is being transmitted to the International Bureau.

This International Search Report consists of a total of 3 sheets.

☒ It is also accompanied by a copy of each prior art document cited in this report.

**1. Basis of the report**

- a. With regard to the **language**, the international search was carried out on the basis of the international application in the language in which it was filed, unless otherwise indicated under this item.

☐ the international search was carried out on the basis of a translation of the international application furnished to this Authority (Rule 23.1(b)).

- b. With regard to any **nucleotide and/or amino acid sequence** disclosed in the international application, the international search was carried out on the basis of the sequence listing :

☐ contained in the international application in written form.

☐ filed together with the international application in computer readable form.

☒ furnished subsequently to this Authority in written form.

☒ furnished subsequently to this Authority in computer readable form.

☒ the statement that the subsequently furnished written sequence listing does not go beyond the disclosure in the international application as filed has been furnished.

☒ the statement that the information recorded in computer readable form is identical to the written sequence listing has been furnished

2. ☐ **Certain claims were found unsearchable** (See Box I).

3. ☐ **Unity of invention is lacking** (see Box II).

**4. With regard to the title,**

☒ the text is approved as submitted by the applicant.

☐ the text has been established by this Authority to read as follows:

**5. With regard to the abstract,**

☒ the text is approved as submitted by the applicant.

☐ the text has been established, according to Rule 38.2(b), by this Authority as it appears in Box III. The applicant may, within one month from the date of mailing of this international search report, submit comments to this Authority.

**6. The figure of the drawings to be published with the abstract is Figure No.**

☐ as suggested by the applicant.

☐ because the applicant failed to suggest a figure.

☐ because this figure better characterizes the invention.

☐ None of the figures.

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/08319

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/47 C12Q1/68 C07K16/18 G01N33/566

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ROUILLAC C ET AL: "Molecular basis of the altered antigenic expression of RhD in weak D (Du) and RhC/e in Rn phenotypes" BLOOD, vol. 87, no. 11, 1 June 1996 (1996-06-01), pages 4853-4861, XP002108679 cited in the application page 4853, last paragraph; table 1 ---	1-3,9, 14-26, 45-48
X	MAAS JH ET AL.: "Rh-D-Genotypisierung auf Exon 2, 5 und 7 von deutschen und japanischen Blutspendern mit sequenzspezifischer Polymerasekettenreaktion" BEITRÄGE ZUR INFUSIONSTHERAPIE UND TRANSFUSIONSMEDIZIN, vol. 34, 1997, pages 203-209, XP002108678 paragraph bridging pages 204 and 205 --- -/-	1-3,9, 14, 17-26,48

☒ Further documents are listed in the continuation of box C.☐ Patent family members are listed in annex.

## \* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&amp;" document member of the same patent family

Date of the actual completion of the international search

9 July 1999

Date of mailing of the international search report

22/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Cupido, M

## INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 98/08319

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category °	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	AVENT ND ET AL: "Evidence of genetic diversity underlying Rh D-, weak D (Du), and partial D phenotypes as determined by multiplex polymerase chain reaction analysis of the RHD gene" BLOOD, vol. 89, no. 7, 1 April 1997 (1997-04-01), pages 2568-2577, XP002108680 cited in the application table 2	1-3, 9-14, 17-26, 48
P, X	--- LEGLER TJ ET AL: "RHD genotyping in weak D phenotypes by multiple polymerase chain reactions" TRANSFUSION, vol. 38, May 1998 (1998-05), pages 434-440, XP002108681 the whole document	1-48
T	--- WAGNER FF ET AL: "Molecular basis of weak D phenotypes" BLOOD, vol. 93, no. 1, 1 January 1999 (1999-01-01), pages 385-393, XP002108682 the whole document -----	1-48



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/12, C07K 14/47, C12Q 1/68,</b> <b>C07K 16/18, G01N 33/566</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/37763</b> <b>(43) International Publication Date:</b> 29 July 1999 (29.07.99)
<b>(21) International Application Number:</b> PCT/EP98/08319 <b>(22) International Filing Date:</b> 18 December 1998 (18.12.98)  <b>(30) Priority Data:</b> 98101203.2                      23 January 1998 (23.01.98)                      EP  <b>(71) Applicant (for all designated States except US):</b> DRK BLUT- SPENDEDIENST BADEN-WÜRTTEMBERG GMBH [DE/DE]; Helmholtzstrasse 10, D-89081 Ulm (DE).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> FLEGEL, Willy, A. [DE/DE]; Grenzstrasse 11, D-64807 Dieburg (DE). WAG- NER, Franz, F. [DE/DE]; Burgunderweg 9, D-89075 Ulm (DE).  <b>(74) Agent:</b> VOSSIUS & PARTNER; P.O. Box 86 07 67, D-81634 Munich (DE).		<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims</i> <i>and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> 16 September 1999 (16.09.99)
<b>(54) Title:</b> NOVEL NUCLEIC ACID MOLECULES CORRELATED WITH THE RHESUS WEAK D PHENOTYPE		
<b>(57) Abstract</b> <p>The present invention relates to novel nucleic acid molecules encoding a Rhesus D antigen contributing to the weak D phenotype which are characterized by one or a combination of missense mutations or by a gene conversion involving exons 6 to 9 of the <i>RHD</i> and <i>RHCE</i> genes. The present invention further relates to vectors comprising the nucleic acid molecules of the invention, to hosts transformed with said vectors, to proteins encoded by said nucleic acid molecules and to methods of producing such polypeptides. The fact that missense mutations and the conversion referred to above can be directly correlated to the weak D phenotype has a significant impact on the routine testing of blood samples. For example, oligonucleotides and antibodies can now be designed that generally allow the detection of weak D phenotypes in a sample. Such oligonucleotides, antibodies as well as a variety of diagnostic methods all fall within the scope of the present invention. RhD antigens encoded by the novel nucleic acid molecules may be used for the characterization, standardization and quality control of monoclonal and polyclonal anti-D antisera. Finally, the invention relates to a kit useful for testing for the presence of weak D phenotypes.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

### Novel nucleic acid molecules correlated with the Rhesus weak D phenotype

The present invention relates to novel nucleic acid molecules encoding a Rhesus D antigen contributing to the weak D phenotype which are characterized by one or a combination of missense mutations or by a gene conversion involving exons 6 to 9 of the *RHD* and *RHCE* genes. The present invention further relates to vectors comprising the nucleic acid molecules of the invention, to hosts transformed with said vectors, to proteins encoded by said nucleic acid molecules and to methods of producing such polypeptides. The fact that missense mutations and the conversion referred to above can be directly correlated to the weak D phenotype has a significant impact on the routine testing of blood samples. For example, oligonucleotides and antibodies can now be designed that generally allow the detection of weak D phenotypes in a sample. Such oligonucleotides, antibodies as well as a variety of diagnostic methods all fall within the scope of the present invention. RhD antigens encoded by the novel nucleic acid molecules may be used for the characterization, standardization and quality control of monoclonal and polyclonal anti-D antisera. Finally, the invention relates to a kit useful for testing for the presence of weak D phenotypes.

The Rhesus D antigen (ISBT 004.001; RH1) carried by the RhD protein is the most important blood group antigen determined by a protein. It is still the leading cause for the hemolytic disease of the newborn (Mollison et al.1993). About 0.2% to 1% of whites have red cells with a reduced expression of the D antigen (weak D, formerly D<sup>u</sup>) (Mourant et al.1976; Stratton, 1946; Wagner et al.1995). A small fraction of weak D samples is explained by qualitatively altered RhD proteins, called partial D (Salmon et al.1984) and frequently caused by *RHD/RHCE* hybrid alleles (recently reviewed in Huang, 1997). Another fraction is caused by the suppressive effects of Cde haplotypes in trans position (Ceppellini et al.1955). These weak D likely possess the normal *RHD* allele, because the carriers' parents and children express often a

normal RhD antigen density. Such weak D show only a minor reduction of RhD antigen expression, were loosely called „high grade D<sup>u</sup>“ and typed today often as normal RhD, because of the increased sensitivity of monoclonal anti-D antibodies.

The majority of moderately to strongly weakened antigen D are due to genotype(s) located either at the Rhesus genes' locus itself or closeby, because the weak D expression is inherited along with the RhD phenotype (Stratton, 1946). Besides the mere quantitative reduction, no qualitative differences could be discerned in the RhD antigen of this most prevalent type of weak D. Two recent studies addressed the molecular cause of the prevalent weak D phenotypes. Both groups, Rouillac et al. (1996) and Beckers et al. (1997), performed RT-PCR and found no mutations when sequencing of *RHD* cDNA in weak D samples. Using semi-quantitative RT-PCR, Rouillac et al. (1996) reported reduced steady-state levels of *RHD* transcripts in weak D samples and disclosed, that their observations provided direct evidence of an only quantitative difference in RhD between normal and weak D red blood cells. In a similar approach, Beckers et al. (1995 and 1997), however, found no differences in the amounts of *RHD* transcripts and excluded an excess of splice variants (Kajii et al.1995), whose products may be inadequately or not at all incorporated in the red cell membrane (Beckers et al.1997). They concluded that weak D is not caused by regulatory defects of the transcription process and proposed unidentified regulatory genes or factors involved in the Rh-related complex as possible causes of weak D. Hence, while the mechanism of weak D expression remained equivocal, no molecular cause was established.

Screening of random weak D samples by PCR for *RHD* specific polymorphisms confirmed PCR amplification patterns representative for a normal *RHD* allele (Avent et al.1997b; Legler et al.1997). However, evidence was accumulating that very few weak D not known to represent partial D, may carry structurally abnormal *RHD* alleles: Four of 44 weak D in England lacked *RHD* specific intron 4 PCR amplicons (Avent et al.1997b) and one out of 94 weak D in Northern Germany lacked *RHD* specific exon 5 PCR amplicons (Legler et al.1997). In the latter sample, the nucleotide T at position 667 was substituted by the *RHCE* specific G coding for a F223V amino acid substitution (TJ Legler and A Humpe, personal communications).



Thus, aberrant alleles were observed only in a small fraction of weak D phenotypes rendering the possibility unlikely that these changes at the molecular level were indeed responsible for the general phenomenon of the weak D phenotype; see also Aubin et al.1997; Avent et al.1997b; Fukumori et el., 1997; Huang, 1997, Issitt and Telen, 1996; Roubinet et al., 1996. Consequently, the combined prior art failed to hitherto provide an conveniently applicable and reliable means to detect the weak Rhesus D phenotype in a sample.

Accordingly, the technical problem underlying the present invention was to establish such a means as well as methods that can conveniently and widely be employed in the analysis of the Rhesus weak D phenotype.

The solution to said technical problem is achieved by providing the embodiments characterized in the claims. Thus, the present invention relates to a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, in its transmembrane and/or intracellular regions.

In accordance with the present invention, the term "contributing to the weak D phenotype" implies an active role of the mutation which may be caused by an amino acid exchange whereas the term "indicative of the weak D phenotype" does not necessarily imply such a role but may also refer to a silent mutation. Such a silent mutation may, for example, occur in conjunction with other mutations such as missense mutations which are addressed in more detail herein below.

In accordance with the present invention, it was found that the observed missense mutations are not only associated with, but truly caused a reduced RhD protein integration into the red blood cells' membranes. Thus, by the present invention it is demonstrated that (i) weak D alleles evolved independently in the different haplotypes, each distinct event being associated with a change in the RhD coding sequence; (ii) no sample occurred with a normal coding sequence despite the

observation of 16 different alleles in 164 samples; and (iii) type and distribution of the observed nucleotide substitutions was not compatible with the null hypotheses of random changes.

The finding that missense mutations in *RHD* led to reduced D antigen expression, fitted into the current model of RhD membrane integration; see Table 7. Both Rh proteins occur in a complex with the Rh50 protein, which can be joined by several additional proteins, like LW, CD47, and glycophorin B (Huang, 1997). The expression of the whole Rh complex depends on the integrity of at least one Rh protein (JP Cartron, oral presentation at the ISBT/DGTI conference, Frankfurt, September 1997) and the Rh50 protein (Cherif-Zahar et al.1996). Subtle structural changes in the Rh50 protein caused by missense mutations are sufficient to prevent the expression of the Rh complex (Cherif-Zahar et al.1996). Likewise, such subtle structural changes in the RhD protein appear to also affect the expression of the Rh complex involving RhD.

Based on the distribution and kind of amino acid substitutions, a general picture of the relationship of RhD structure and RhD expression can now be established: All amino acid substitutions in weak D are located in the intracellular or transmembrane parts of the RhD protein where the alignment was carried out in accordance with the above mentioned current model (see Table 7). Known *RHD* alleles with exofacial substitutions (Avent et al.1997a; Jones et al.1997; Liu et al.1996; Rouillac et al.1995) were discovered by virtue of their partial D antigen, but may display discrete ( $D^{nu}$  and  $D^{vi}$ ) to moderate ( $D''$ , DHR and DHMi) reductions in RhD expression (Flegel and Wagner, 1996; Jones et al.1997; Jones et al.1996). Most substitutions reported in accordance with this invention were nonconservative and the introduced amino acids, in particular proline, likely disrupted the secondary or tertiary structure. Two weak D alleles (type 2 and 11) were associated with conservative substitutions indicating that the involved amino acid regions at positions 295 and 385 were very important for an optimal RhD membrane integration. In two alleles (type 4 and type 14), parts of exon 4 and 5 were substituted by the corresponding parts of the *RHCE* gene. Similar exchanges occurred in  $D^{vi}$  type I and  $D^{vi}$  type II that exhibited a considerably reduced RhD protein expression (Jones et al.1996), too. Previous paradoxical observations

can be explained, if the N152T substitution in exon 3 is considered to facilitate the membrane integration: (i) D<sup>IIIa</sup> (Huang et al.1997), differing from weak D type 4 by the N152T substitution only, has a normal RhD antigen density (Jones et al.1996), and (ii) D<sup>IIIc</sup>, D<sup>IVa</sup>, and D<sup>VI</sup> type III harbouring the N152T substitution have enhanced antigen densities (Flegel et al.1997; Jones et al.1996) compared to their appropriate controls (normal RhD and D<sup>VI</sup> type II).

Several phenotypes with weak D expression, like D<sup>VI</sup>, D<sup>V</sup>, DBT, some D<sup>IV</sup> and DFR, were recognized long ago as separate entities by their carriers' propensity to produce anti-D (Lomas et al.1994; Tippett and Sanger, 1977; Tippett and Sanger, 1962). These phenotypes were subsequently confirmed and grouped by distinct reaction patterns with monoclonal anti-D (Lomas et al.1993; Lomas et al.1989; Scott, 1996). A serologic classification of most weak D phenotypes, however, has not been successful, because they lacked a consistent reaction pattern with monoclonal anti-D and their carriers seemed not prone to anti-D immunization (Moore, 1984). There was even no defined borderline between normal D and weak D (Agre et al.1992; Moore, 1984; Nelson et al.1995). Nevertheless, variability of the RhD antigen density (antigens per cell) in weak D phenotypes (Hasekura et al.1990; Jones et al.1996; Nelson et al.1995; Nicholson et al.1991; Tazzari et al.1994; Wagner, 1994) and rare aberrant patterns in *RHD* PCR (Avent et al.1997b; Legler et al.1997) did not exclude an underlying molecular diversity. The present invention for the first time allows for the convenient classification of weak D and for the unambiguous correlation of distinct alleles with clinical data. In conjunction with previously defined rare *RHD* alleles, the exact molecular definition of most phenotypes with reduced D antigen density has now become possible. In the case that patients carrying particular molecular types of weak D were prone to develop anti-D, the classification made possible by the present invention will help to guide a Rhesus negative transfusion policy. The availability of weak D samples that are characterized in regard to molecular structure and RhD antigen densities will promote the quality assurance of anti-D reagents. They should reliably type probands as RhD positive, whose RhD proteins are not prone to frequent anti-D immunization (Wagner et al.1995). Therefore, the use of RhD negative red blood cell units for transfusions to weak D patients, which has been justified by a

presumed potential for anti-D immunization, can finally be reduced to a minimum, which can be scientifically deduced.

Additionally, it was found in accordance with the invention that the mutations cluster in certain stretches of the Rhesus D polypeptide. Further, a gene conversion correlating with the weak D phenotype was detected. Thus, the invention also relates to a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule

- (a) carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, in amino acid positions 2-16, 114-149, 179-225 or/and 267 to 397 with the proviso that said D antigen does not carry a single missense mutation leading to a substitution of phenylalanine in amino acid position 223 by valine or of threonine in position 283 by isoleucine; or
- (b) carrying a gene conversion involving exons 6 to 9 which are replaced by the corresponding exons of the *RHCE* gene.

All the missense mutations found in accordance with the present invention and located in the above regions are associated with the transmembrane region or the intracellular portion of the polypeptide when the above indicated current model of RhD is employed. However, when different models are employed, the mutations associated with the weak D phenotype may also be found in the extracellular regions. The above regions also comprise amino acid positions which are located in the extracellular regions when the current model is applied. Said positions might also be mutated and correlatable with the weak D phenotype. Such mutations also fall within the scope of the application.

In addition to the missense mutations, a gene conversion indicative of weak D was identified. Said conversion can be used for diagnostic purposes basically to the same extent as the missense mutations. In accordance with the invention, the breakpoints are determined to be in introns 5 and 9; see also Fig. 3.

The mutants referred to above and further throughout this specification can be conveniently employed for the characterization of monoclonal and polyclonal

antibodies used in connection with RhD diagnosis, prophylaxis and treatment. For example, by expressing desired nucleic acid molecules encoding such mutants in a suitable system, reactivity profiles of said antibodies or antisera can be established. The mutants can also be employed for the characterization of monoclonal and polyclonal antibodies that are used as secondary antibodies, for example, anti-globulin and anti-human-globulin antisera.

Preferably, the missense mutation causes an amino acid substitution in position 3, 10, 16, 114, 149, 182, 198, 201, 220, 223, 270, 276, 277, 282, 294, 295, 307, 339, 385 or 393 or a combination of/or involving said substitutions.

This preferred embodiment, besides the single mutations indicated, may comprise a combination of these substitutions. Additionally, it contemplates the possibility that one or more of said substitutions are involved, and additional mutations such as mutations leading to substitutions are present. In accordance with the present invention, it is understood that such additional mutations may be tested for when assessing RhD status in a sample. A finding of such a mutation will allow the person skilled in the art to conclude that other mutations identified in this specification occurring in combination with said first mutation will be present. Accordingly, such embodiments reflecting the detection of additional mutations occurring in combination with the mutations identified in this specification are also comprised by the invention.

In a particularly preferred embodiment of the nucleic acid molecule of the invention, said amino acid substitution in position 3 is from Ser to Cys, in position 10 from Arg to Gln, in position 16 from Trp to Cys, in position 114 from Arg to Trp, in position 149 from Ala to Asp, in position 182 from Ser to Thr, in position 198 from Lys to Asn, in position 201 from Thr to Arg, in position 220 from Trp to Arg, in position 223 from Phe to Val, in position 270 from Val to Gly, in position 276 from Ala to Pro, in position 277 from Gly to Glu, in position 282 from Gly to Asp, in position 294 from Ala to Pro, in position 295 from Met to Ile, in position 307 from Gly to Arg, in position 339 from Gly to Glu, in position 385 from Gly to Ala and in position 393 from Trp to Arg.

In a further preferred embodiment of the nucleic acid molecule of the invention, said missense mutation occurs in nucleotide position 8, 29, 48, 340, 446, 544, 594, 602, 658, 667, 809, 819, 826, 830, 845, 880, 885, 919, 1016, 1154 and 1177 or in a combination of said positions.

Particularly preferred is that said missense mutation in position 8 is from C to G, in position 29 from G to A, in position 48 from G to C, in position 340 from C to T, in position 446 from C to A, in position 544 from T to A, in position 594 from A to T, in position 602 from C to G, in position 658 from T to C, in position 667 from T to G, in position 809 from T to G, in position 819 from G to A, in position 826 from G to C, in position 830 from G to A, in position 845 from G to A, in position 880 from G to C, in position 885 from G to T, in position 919 from G to A, in position 1016 from G to A, in position 1154 from G to C and in position 1177 from T to C.

In the case that combinations of missense mutations are involved in the generation of weak D phenotypes, it is preferred that said combination of substitutions is in positions 182, 198 and 201 and is preferably S182T, K198N, T201R or in position 201 and 223 and is preferably T201R and F223V, or in position 16, 201 and 223 and is preferably W16C, T201R and F223V.

Most preferably, said combination of missense mutations comprises positions 544, 594 and 602 and is preferably T→A at position 544, A→T at position 594 and C→G at position 602 or comprises positions 602, 667 and 819 and is preferably C→G at position 602, T→G at position 667 and G→A at position 819, or comprises positions 48, 602, 667 and 819 and is preferably G→C at position 48, C→G at position 602, T→G at position 667 and G→A at position 819.

Although the nucleic acid molecule of the invention may be of various origin including (semi) synthetic origin, it is preferred that the nucleic acid molecule is mRNA or genomic DNA. Standard procedures may be employed to obtain any of the above nucleic acids; see, for example, Sambrook, et al., "Molecular Cloning, A Laboratory Manual", 2<sup>nd</sup> ed. 1989, CSH Press, Cold Spring Harbor, N.Y.

The invention also relates to a vector comprising the nucleic acid molecule of the invention.

The vector may be used for propagation and/or expression or may be designed for gene transfer or targeting purposes. Methods of producing such vectors are well known in the art. The same holds true for cloning the nucleic acids of the mutation into said vectors, as well as the propagation of vectors in suitable hosts, etc.

The vector may particularly be a plasmid, a cosmid, a virus or a bacteriophage used conventionally in genetic engineering that comprise the nucleic acid molecule of the invention. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the nucleic acid molecules or vector of the invention into targeted cell populations. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook, *Molecular Cloning A Laboratory Manual*, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel, *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y. (1989). Alternatively, the polynucleotides and vectors of the invention can be reconstituted into liposomes for delivery to target cells. The vectors containing the nucleic acid molecules of the invention can be transferred into the host cell by well-known methods, which vary depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment or electroporation may be used for other cellular hosts; see Sambrook, *supra*.

Such vectors may comprise further genes such as marker genes which allow for the selection of said vector in a suitable host cell and under suitable conditions. Preferably, the nucleic acid molecule of the invention is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic cells. Expression of said polynucleotide comprises transcription of the polynucleotide into a translatable mRNA. Regulatory elements ensuring expression in eukaryotic cells, preferably mammalian cells, are well known to those skilled in the art. They usually comprise regulatory sequences ensuring initiation of transcription and optionally poly-A signals ensuring termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational

enhancers, and/or naturally-associated or heterologous promoter regions. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the PL, lac, trp or tac promoter in *E. coli*, and examples for regulatory elements permitting expression in eukaryotic host cells are the AOX1 or GAL1 promoter in yeast or the CMV-, SV40-, RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the nucleic acid molecule. Furthermore, depending on the expression system used leader sequences capable of directing the polypeptide to a cellular compartment or secreting it into the medium may be added to the coding sequence of the polynucleotide of the invention and are well known in the art. The leader sequence(s) is (are) assembled in appropriate phase with translation, initiation and termination sequences, and preferably, a leader sequence capable of directing secretion of translated protein, or a portion thereof, into the periplasmic space or extracellular medium. Optionally, the heterologous sequence can encode a fusion protein including an C- or N-terminal identification peptide imparting desired characteristics, e.g., stabilization or simplified purification of expressed recombinant product. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (In-vitrogene), or pSPORT1 (GIBCO BRL).

Preferably, the expression control sequences will be eukaryotic promoter systems in vectors capable of transforming or transfecting eukaryotic host cells, but control sequences for prokaryotic hosts may also be used.

As mentioned above, the vector of the present invention may also be a gene transfer or targeting vector. Gene therapy, which is based on introducing therapeutic genes into cells by ex-vivo or in-vivo techniques is one of the most important applications of gene transfer. Suitable vectors and methods for in-vitro or in-vivo gene therapy are described in the literature and are known to the person skilled in the art; see, e.g., Giordano, *Nature Medicine* 2 (1996), 534-539; Schaper, *Circ. Res.* 79 (1996), 911-919; Anderson, *Science* 256 (1992), 808-813; Isner, *Lancet* 348 (1996), 370-374; Muhlhauser, *Circ. Res.* 77 (1995), 1077-1086; Wang, *Nature Medicine* 2 (1996), 714-716; WO94/29469; WO 97/00957 or Schaper, *Current Opinion in*



Biotechnology 7 (1996), 635-640, and references cited therein. The polynucleotides and vectors of the invention may be designed for direct introduction or for introduction via liposomes, or viral vectors (e.g. adenoviral, retroviral) into the cell. Preferably, said cell is a germ line cell, embryonic cell, or egg cell or derived therefrom, most preferably said cell is a stem cell.

Additionally, the invention relates to a host transformed with the vector of the invention.

Appropriate hosts comprise transgenic animals, cells such as bacteria, yeast cells, animal, preferably mammalian cells, fungal cells or insect cells. Transformation protocols including transfection, microinjection, electroporation, etc., are also well known in the art.

Further, the invention relates to a method of producing a Rhesus D antigen contributing to the weak D phenotype comprising culturing the host of the invention under suitable conditions and isolating the Rhesus D antigen produced.

It is preferred that the antigen is exported into the culture medium where it can be collected according to conventions/methods. The term "culturing" as used in accordance with the present invention also comprises the raising of transgenic animals. Using appropriate vectors constructions and optionally appropriate feeds, the antigen may, e.g., be isolated from milk of, e.g. transgenic cows.

The invention additionally relates to Rhesus D antigen encoded by the nucleic acid molecule of the invention or produced by the method of the invention.

Preferably, the antigen is in the same way post transitionally modified and has the same chemical structure as naturally occurring antigen. Accordingly, said antigen, when produced by the method of the invention, is preferably produced in human cells.

Furthermore, the invention relates to an oligonucleotide hybridizing under stringent conditions to a portion of the nucleic acid molecule of the invention comprising said at least one missense mutation or to the complementary portion

thereof or hybridizing to a breakpoint of the gene conversion identified here in the above.

In this embodiment of the invention, it is understood that the oligonucleotides hybridizes directly to the mutated sequence or to the breakpoint. The setting of stringent hybridization conditions is well described, for example, in Sambrook et al, "Molecular Cloning, A Laboratory Handbook" CSH Press, Cold Spring Harbor 1989 or Hames and Higgins, "Nucleic acid hybridization, a practical approach", IRL Press, Oxford (1985). Thus, the detection of the specifically hybridizing sequences will usually require hybridization and washing conditions such as 0.1xSSC, 0.1% SDS at 65°. As is well known, the length of the probe and the composition of the nucleic acid to be determined constitute further parameters of the stringent hybridization conditions. Preferably, the oligonucleotide is a deoxynucleotide. It is further preferred that the oligonucleotide comprises 12 to 50 nucleotides and more preferably 15 to 24 nucleotides. The hybridization to the breakpoint may be under stringent or non-stringent conditions. An example of non-stringent hybridization conditions is hybridization and washing at 50°C in 4xSSC, 0.1% SDS.

Further, the invention relates to an antibody or an aptamer specifically binding to the Rhesus D antigen of the invention.

The antibody may be tested and used in any serologic technique well known in the art, like agglutination techniques in tubes, gels, solid phase and capture techniques with or without secondary antibodies, or in flow cytometry with or without immunofluorescence enhancement.

The antibody of the invention may be a monoclonal antibody or an antibody derived from or comprised in a polyclonal antiserum. The term "antibody", as used in accordance with the present invention, further comprises fragments of said antibody such as Fab, F(ab')<sub>2</sub>, Fv or scFv fragments; see, for example, Harlow and Lane, "Antibodies, A Laboratory Manual" CSH Press 1988, Cold Spring Harbor, N.Y. The antibody or the fragment thereof may be of natural origin or may be (semi) synthetically produced. Such synthetic products also comprises non-proteinaceous as semi-proteinaceous material that has the same or essentially the same binding

specificity as the antibody of the invention. Such products may, for example, be obtained by peptidomimetics.

Additionally, the invention relates to an antibody or an aptamer or a phage specifically binding to the wild type Rhesus D antigen or to aberrant D Rhesus antigens but not to the Rhesus D antigen of the invention. The antibody may be tested and used in any serologic technique well known in the art, like agglutination techniques in tubes, gels and solid phase techniques, capture techniques or flow cytometry with immunofluorescence.

As regards, the definition, testing and origin of the antibody or the aptamer, the same definitions as above apply here.

As regards the term "aberrant Rhesus D antigen", the term comprises prior art missense mutations as well as prior art conversions found in RHD genes and the corresponding antigens.

The term "aptamer" is well known in the art and defined, e.g., in Osborne et al., Curr. Opin. Chem. Biol. 1 (1997), 5-9 or in Stall and Szoka, Pharm. Res. 12 (1995), 465-483.

Furthermore, the invention relates to a method for testing for the presence of a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising hybridizing the oligonucleotide of the invention or an oligonucleotide hybridizing to a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T under stringent conditions to nucleic acid molecules comprised in the sample obtained from a human and detecting said hybridization.

Preferably, the method of the invention further comprises digesting the product of said hybridization with a restriction endonuclease or subjecting the product of said hybridization to digestion with a restriction endonuclease and analyzing the product of said digestion.

This preferred embodiment of the invention allows by convenient means, the differentiation between an effective hybridization and a non-effective hybridization. For example, if the wild type Rhesus D antigen comprises an endonuclease restriction site, the hybridized product will be cleavable by an appropriate restriction enzyme whereas a mutated sequence will yield no double-stranded product or will not comprise the recognizable restriction site and, accordingly, will not be cleaved. Alternatively, the hybridizing oligonucleotide may only hybridize to the mutated sequence. In this case, only a hybrid comprising the mutated sequence, but not the wild type sequence, will be cleaved by the appropriate restriction enzyme. The analysis of the digestion product can be effected by conventional means, such as by gel electrophoresis which may be optionally combined by the staining of the nucleic acid with, for example, ethidium bromide. Combinations with further techniques such as Southern blotting are also envisaged.

Detection of said hybridization may be effected, for example, by an anti-DNA double-strand antibody or by employing a labeled oligonucleotide. Conveniently, the method of the invention is employed together with blotting techniques such as Southern or Northern blotting and related techniques. Labeling may be effected, for example, by standard protocols and includes labeling with radioactive markers, fluorescent, phosphorescent, chemiluminescent, enzymatic labels, etc.

The invention additionally relates to a method of testing for the presence of a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising determining the nucleic acid sequence of at least a portion of the nucleic acid molecule of the invention, said portion encoding at least one of said missense mutations or a breakpoint of said gene conversion or a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense

mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T.

Preferably, the method of the invention further comprises, prior to determining said nucleic acid sequence, amplification of at least said portion of said nucleic acid molecule.

Preferably, amplification is effected by polymerase chain reaction (PCR). Other amplification methods such as ligase chain reaction may also be employed.

Furthermore, the invention relates to a method for testing for the presence of a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising carrying out an amplification reaction wherein at least one of the primers employed in said amplification reaction is the oligonucleotide of the invention or an oligonucleotide hybridizing to a nucleic acid molecule encoding a Rhesus D antigen contributing to the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T and assaying for an amplification product.

The method of the invention will result in an amplification of only the target sequence, if said target sequence carries the or at least one mutation. This is because the oligonucleotide will under preferably stringent hybridization conditions not hybridize to the wild type sequence (with the consequence that no amplification product is obtained) but only to the mutated sequence. Naturally, primer oligonucleotides hybridizing to one or more as one, such as two mutated sequences

may be employed in the method of the invention. The latter embodiment may be favorable in cases where combinations of mutations are tested for. It is important to note that not all or none of said mutations are necessarily missense mutations. This may be true for cases where other types of mutations occur in combination with the above missense mutations or with the above gene conversion.

Preferably, in the method of the invention said amplification or amplification reaction is or is effected by the polymerase chain reaction (PCR). Other amplification methods such as ligase chain reaction may also be employed.

Further, the invention relates to a method for testing for the presence of a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising assaying a sample obtained from a human for specific binding to the antibody or aptamer or phage of the invention or to an antibody or aptamer or phage to a Rhesus D antigen contributing to or indicative of the weak D phenotype and encoded by a nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T.

Testing for binding may, again, involve the employment of standard techniques such as ELISAs; see, for example, Harlow and Lane, "Antibodies, A Laboratory Manual" CSH Press 1988, Cold Spring Harbor.

The invention also relates to the method of testing a sample for the presence of wild type Rhesus D antigen and the absence of the Rhesus D antigen of the invention comprising assaying a sample obtained from a human for specific binding to the antibody or aptamer or phage of the invention, said antibody or aptamer or phage specifically binding to the wild type Rhesus D antigen or to aberrant D Rhesus antigens but not to the Rhesus D antigen of the invention.

Results obtained in accordance with their method of invention may well be employed in strategies of blood transfusion, as outlined herein above.

Preferably, in the method of the invention said sample is blood, serum, plasma, fetal tissue, saliva, urine, mucosal tissue, mucus, vaginal tissue, fetal tissue obtained from the vagina, skin, hair, hair follicle or another human tissue.

Furthermore, the method of the invention preferably comprises the step of enrichment of fetal cells. This enrichment may be achieved by using appropriate antibodies, lectins or other reagents specifically binding fetal cells or by any technique attempting the differential separation of maternal and fetal cells, like by density gradients. Also preferably, in said method fetal DNA or mRNA from material tissue like peripheral blood, serum or plasma may be extracted, advantageously according to conventional procedures.

In an additional preferred embodiment of the method of the invention, said nucleic acid molecule or proteinaceous material from said sample is fixed to a solid support.

Preferably, said solid support is a chip.

The advantages of chips are well known in the art and need not be discussed herein in detail. These include the small size as well as an easy access of computer based analysis of analytes.

Furthermore, the present invention relates to the use of the nucleic acid molecule of the invention or of a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T or of a combination thereof for the analysis of a weak Rhesus D phenotype.

The analysis can be effected, for example, on the basis of the methods described herein above.

The invention also relates to the use of the nucleic acid molecule of the invention, the vector of the invention or the Rhesus D antigen of the invention for the assessment of the affinity, avidity and/or reactivity of monoclonal anti-D antibodies or of polyclonal anti-D antisera or of anti-globulin or of anti-human-globulin antisera or of preparations thereof.

The invention also relates to the use of cells, preferably red blood cells, from probands for the assessment of the affinity, avidity and/or reactivity of monoclonal anti-D antibodies or of polyclonal anti-D antisera or of anti-globulin or of anti-human-globulin antisera or of preparations thereof.

Said preparations can be provided according to techniques well known in the art. Said preparations may comprise stabilisators such as albumins, further sodium azide, salt ions, buffers etc. The formulation of the preparation may have an influence on the binding characteristics of the antibodies, as is well known in the art.

For example, in a first step, the Rhesus D gene of a carrier or of a blood donor and its allelic status is analyzed and it is determined whether said gene comprises a mutation that was found in accordance with the present invention. In a second step, said mutation is correlated to a certain RhD antigen density on the surface of red blood cells. Conveniently, said correlation can be established by data provided in the present invention (such as mutations per se) and techniques that are well known in the art (see, e.g. Jones et al. 1996, Flegel and Wagner, 1996). In a third step, the features of an antibody or an antiserum such as reactivity, sensitivity, affinity, avidity, and/or specificity are determined with suitable blood group serological techniques preferably using red blood cells that were molecularly and with respect to the RhD antigen surface density characterized as described in step 2. Such data can be used, for example, in quality controls, standardization, etc.

The invention will be most useful for the characterization, standardization and quality control of monoclonal and polyclonal antisera, preferably anti-D monoclonals or antisera. Further, for example, anti-globulin and anti-human-globulin antisera can



be characterized on the basis of the teachings of the present invention. An appropriately characterized anti-D monoclonal antibody can be conveniently used in RhD diagnostics. For example, a suitably characterized monoclonal antibody will be useful in determining the weak D antigen density on the surface of blood cells. Cut-off values for monoclonal antibodies useful in diagnosis can thus be established. This is important for the quality control of antibodies used in RhD diagnosis.

Thus, the invention also relates to a method for the characterization of monoclonal antibodies or polyclonal antisera or of a preparation thereof, said method comprising

- (a) testing the nucleic acid of sample of a proband for the presence of a mutation as defined in accordance with the invention;
- (b) correlating, on the basis of the mutation status and the allelic status of the *RHD* gene, the nucleic acid with the RhD antigen density on the surface of red blood cells of said proband;
- (c) reacting said monoclonal antibodies or polyclonal antisera or said preparation thereof with a cell carrying the RhD antigen on its surface;
- (d) characterizing said monoclonal antibodies or polyclonal antisera or said preparation thereof on the basis of the results obtain in step (c).

As regards the term "allelic status", this term describes the possibilities that the *RHD* alleles in a proband are present in a homozygous, heterozygous or hemizygous state. Also comprised by this term is the possibility that the two alleles carry two different mutations (including the conversion) defined herein above.

In a preferred embodiment of the method of the invention, said characterization comprises the determination of reactivity, sensitivity, avidity, affinity, specificity and/or other characteristics of antibodies and antisera.

Furthermore preferred is a method wherein said cell carrying the RhD antigen on its surface is a red blood cell.

The invention also relates to a method for determining whether a patient in need of a blood transfusion is to be transfused with Rh D negative blood from a donor comprising the step of testing a sample from said patient for the presence of one or more Rh D antigens of the invention, wherein a positive testing for at least one of said antigens is indicative of the need for a transfusion with Rh negative blood. The invention has important implications for devising a transfusion therapy in humans. For example, it can now be conveniently tested whether the patient actually needs a transfusion with a Rh D negative blood or whether such precautions need not be taken.

The transfusion of red blood cells of some molecularly defined subgroups of the weak D phenotype determined by such methods may be immunogenic, if carriers of the wild type Rhesus D antigen, an aberrant D antigen or another weak D type of the invention were transfused by some subgroup of the weak D phenotype. Such carriers, like blood donors, may be determined by previously established methods in the art or by methods established in the invention and subsequently the transfusion of some subgroups of the weak D phenotype may be avoided.

Furthermore, the invention relates to a method for determining whether blood of a donor may be used for transfusion to a patient in need thereof comprising the step of testing a sample from said donor for the presence of one or more Rh D antigens of the invention, wherein a positive testing for at least one of said antigens excludes the transfusion of patients that are typed as having wild type Rh D antigen or (a) weak D type(s) other than the weak D type(s) of said donors.

On the basis of the method of the invention, it is advantageous and desired to avoid transfusion of a patient with weak D typed blood from a donor, if the weak D antigens in both donor and recipient are not totally identical.

The samples referred to in the above recited methods may be samples that are referred to throughout the specification, such as blood, serum, etc.

As regards the guidelines for transfusing a patient on the basis of any of the above recited methods, the utmost care must be taken that suboptimal transfusion policy is avoided. The risk factor is always to be considered by the physician in charge. In all cases, the potential risk for the patient is to be minimized.

The present invention is particularly suitable for establishing criteria which will guide the future strategies in blood transfusion policy. According to the molecular criteria established by the invention, the weak D phenotype can be grouped. Some molecularly defined subgroups of the weak D phenotype determined by such methods may be prone to immunization, if the carriers were transfused with the wild type Rhesus D antigen, an aberrant D antigen or another weak D type of the invention, and may produce an anti-D. Such carriers may be determined by methods established in the invention and subsequently transfused with Rhesus negative blood components, like erythrocyte, thrombocyte and plasma blood units. The majority of carriers with weak D phenotype is by the current art not considered prone to be immunized in such a way by Rhesus D positive blood transfusions and may, hence, by the means established by the invention safely be transfused Rhesus D positive, because of their classification to a distinct weak D type according to the present invention.

The invention also relates to the use of a phage, aptamer, monoclonal antibody or a polyclonal antisera or a preparation thereof as characterized in the present invention for RhD antigen determination.

In a preferred embodiment of said use, said RhD antigen determination is effected in connection with blood group typing.

Furthermore, the invention relates to a preparation comprising the antibody or aptamer or phage of the invention.

The weak D types defined by the invention correlate with certain RhD epitope and RhD antigen densities, i.e. RhD antigens per cell expressed on the red blood cell surface (Flegel and Wagner 1996) (data from few examples are provided in Table 8) Antibodies and preparations thereof may be tested by any standard blood group serology technique with one or more weak D types of the invention. The reactivity,

sensitivity, avidity, affinity, specificity and/or other characteristics of antibodies and antisera known in the art may be determined by its reaction with one or more weak D types of the invention under predetermined conditions in standard blood group serology techniques well known in the art. The preparation may be a diagnostic or pharmaceutical preparation.

The pharmaceutical composition of the present invention may further comprise a pharmaceutically acceptable carrier and/or diluent. Examples of suitable pharmaceutical carriers are well known in the art and include phosphate buffered saline solutions, water, emulsions, such as oil/water emulsions, various types of wetting agents, sterile solutions etc. Compositions comprising such carriers can be formulated by well known conventional methods. These pharmaceutical compositions can be administered to the subject at a suitable dose. Administration of the suitable compositions may be effected by different ways, e.g., by intravenous, intraperitoneal, subcutaneous, intramuscular, topical, intradermal, intranasal or intrabronchial administration. The dosage regimen will be determined by the attending physician and clinical factors. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. A typical dose can be, for example, in the range of 0.001 to 1000  $\mu\text{g}$ ; however, doses below or above this exemplary range are envisioned, especially considering the aforementioned factors. Generally, the regimen as a regular administration of the pharmaceutical composition should be in the range of 1  $\mu\text{g}$  to 10 mg units per day. If the regimen is a continuous infusion, it should also be in the range of 1  $\mu\text{g}$  to 10 mg units per kilogram of body weight per minute, respectively. Progress can be monitored by periodic assessment. The compositions of the invention may be administered locally or systemically. Administration will generally be parenterally, e.g., intravenously; DNA may also be administered directly to the target site, e.g., by biolistic delivery to an internal or external target site or by catheter to a site in an artery. Preparations for parenteral administration include sterile aqueous or non-aqueous solutions, suspensions, and emulsions. Examples of non-aqueous solvents are propylene glycol, polyethylene glycol, vegetable oils such as olive oil, and injectable organic esters such as ethyl

oleate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's, or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be present such as, for example, antimicrobials, anti-oxidants, chelating agents, and inert gases and the like. Furthermore, the pharmaceutical composition of the invention may comprise further agents such as interleukins or interferons depending on the intended use of the pharmaceutical composition.

An antibody and its preparation may be characterized by its reaction or lack of reaction to surfaces with certain RhD epitope densities. For example, antibody preparations may be characterized by agglutinating red blood cells with 1,000 RhD antigens per cell - a RhD antigen density deliberately chosen to be met for quality control purposes.

The invention also relates to treating a pregnant woman being Rhesus D negative or being hemizygous for a mutation defined herein above wherein the child is Rhesus D positive or carries a different mutation defined herein above in a hemizygous state comprising administering anti-D to said woman.

Pregnant women may be currently treated with an anti-D prophylaxis, when a Rhesus negative mother carries a RhD positive fetus. The invention allows the discrimination of an anti-D prophylaxis requirement depending on the status of the mother's and/or the fetus' possessing a RhD protein of the invention. One or more of the RhD proteins of the invention may be prone to immunization of their carriers and, hence, would be indicative for the therapy of the mother. Similarly, one or more RhD proteins of the invention, when carried by the fetus, may be known to be of low immunogenicity to the mother and, hence, would be indicative for the omission of anti-D prophylaxis in difference to current clinical therapy.

The administration can be effected by standard routes and doses which can be defined by the attending physician; Mollison, 1993. Preferably, a monoclonal anti-D or combinations/mixtures of monoclonal anti-Ds is/are administered in doses of 50 µg to

or exceeding 500  $\mu\text{g}$  anti-D antibody/antisera for intravenous or intramuscular administration (Bowman, 1998). For the quality control of these anti-D antibodies/antisera, the results and methods provided by the present invention may be advantageously employed.

The present invention also relates to a method of identifying an antibody  $V_H$  or  $V_L$  chain or a combination thereof or an aptamer specifically binding to a weak D polypeptide of the invention comprising

- (a) contacting the weak D polypeptide of the invention with a phage library displaying  $V_H$  or  $V_L$  chains or combinations thereof on the surface of the phage or with aptamers;
- (b) identifying phage or aptamers that bind to said weak D polypeptide; and optionally
- (c) repeating steps (a) and (b) one or more times.

The preparation of phage library and the screening/identification of desired antibody (chains) per se is well known in the art and reviewed, for example, in Winter et al., *Annu. Rev. Immunol.* 12 (1994), 433-455 and references cited therein. Also, aptamers can be prepared and cloned in phage according to conventional protocols. Whereas single  $V_H$  or  $V_L$  chains may be identified by the method of the invention as binding to the weak D polypeptide of the invention, it is preferred to identify  $V_H$ - $V_L$  combinations expressed by the phage because this situation resembles the situation of natural antibody binding. By repeating steps (a) and (b) one or more times, better binding specificities may be identified. Protocols for the optimization of binding properties such as affinities, including elution steps for removing bound phage, are well established in the art. For example, once a  $V_H$  chain with a convenient binding capacity has been found,  $V_L$  chains may be identified that significantly improve the binding capacity of the antibody, e.g. by replacing the  $V_L$  chain that was associated with the  $V_H$  chain in the first selection step with a more suitable  $V_L$  chain.

The invention also relates to a method of identifying a monoclonal antibody specifically binding to a weak D polypeptide/antigen of the invention comprising

- (a) contacting the weak D polypeptide of the invention with one or more monoclonal antibodies;
- (b) identifying monoclonal antibodies that bind to said weak D polypeptide; and optionally
- (c) repeating steps (a) and (b) one or more times.

The invention also relates to a method of identifying an antibody  $V_H$  or  $V_L$  chain or a combination thereof or an aptamer specifically binding to a weak D polypeptide/antigen of the invention comprising

- (a) contacting the weak D polypeptide and
  - (aa) a second or more weak D weak D polypeptide(s) and/or
  - (ab) a normal Rhesus D polypeptide

wherein the second or more weak D polypeptide(s) and/or the normal Rhesus D polypeptide are present in a molar mass that is higher, equal or less than the weak D polypeptide of (a) with a phage library displaying  $V_H$  or  $V_L$  chains or combinations thereof on the surface of the phage or with aptamers;

- (b) identifying phage or aptamers that bind to said weak D polypeptide of (a); and optionally
- (c) repeating steps (a) and (b) one or more times.

Particularly preferred in step (ab) is that the molar mass of the second weak D polypeptide and the normal Rhesus D polypeptide is higher than that of the weak D polypeptide of (a).

In the case that only one round of selection is employed for the identification (i.e. when step (c) does not apply), it is preferred that the number of weak D polypeptide molecules of (a) is in molar excess over the number of phage particles. The preferred embodiments of the method of identifying an antibody  $V_H$  or  $V_L$  chain or of a combination thereof or of an aptamer described hereinbefore equally apply to this embodiment of the invention.

The invention also relates to a method of identifying a monoclonal antibody specifically binding to a weak D polypeptide/antigen of the invention comprising

- (a) contacting the weak D polypeptide and

- (aa) a second or more weak D polypeptide(s) and/or
  - (ab) a normal D polypeptide
- wherein the second or more weak D polypeptide(s) and/or the normal D polypeptide are present in a molar mass that is higher, equal or less than the weak D polypeptide of (a) with one or more monoclonal antibodies;
- (b) identifying monoclonal antibodies that bind to said weak D polypeptide of (a); and optionally
  - (c) repeating steps (a) and (b) one or more times.

Preferably, the weak D polypeptide is exposed on the surface of a cell. An appropriate surface is the surface of an erythrocyte. However, other host cells may be transfected with a vector suitable for expression of the weak D polypeptide and express the same on their surface. Antibodies may also bind to recombinant proteins or parts of proteins of weak D and purified proteins.

It is further preferred that the polypeptide or host cell is affixed to a solid support. Suitable examples for solid supports are microtiter plates or beads.

In an additionally preferred antibody, subsequent to step (b) or (c), the following step is carried out:

- (d) identifying the amino acid sequence of the  $V_H$  or  $V_L$  chains and/or identifying the nucleic acid sequences encoding said amino acid sequence.

The identification of the amino acid/nucleic acid sequences can be effected according to conventional protocols; see, e.g., Sambrook et al., loc. cit.

Finally, the invention relates to a kit comprising

- (a) the oligonucleotide of the invention; and/or
- (b) the antibody of the invention;
- (c) the aptamer of the invention; and/or
- (d) the phage of the invention.



The kit of the invention which may comprise various types of antibodies described herein above, is particularly suitable for the analysis of weak Ds in samples obtained from humans. The components of the kit may be packaged as appropriate. Preferably, different components are packaged in different vials.

The disclosure content of the documents as cited in this specification is herewith incorporated by reference.

The figures show

**Figure 1.** Schematic representation of the amino acid variations observed in weak D types with single missense mutations. The affected amino acids of the prevalent normal RhD protein and their positions are shown on top. Their substitutions occurring in the weak D types are shown below the bar.

**Figure 2.** The cDNA nucleotide and predicted amino acid sequences of the prevalent allele of the *RHD* gene. The consensus sequences are shown that are deposited in the EMBL nucleotide sequence data base under the accession number X54534 by Avent et al. and modified as noted in the description (C at 1,036). The positions of the nucleotides and amino acids are indicated by the numbers above and below the sequences, respectively.

**Figure 3.** Part of intron 5 of the *RHCE* and *RHD* genes. The nucleotide sequence of the *RHCE* gene is shown. Numbers indicate the position relative to the first base of exon 5 in the *RHCE* gene. Dashes denote nucleotides in the *RHD* gene that are identical to the *RHCE* gene. The 5' breakpoint region (178 bp) of the gene conversion characteristic for D category IV type III is indicated by asterisks. The full intron 5 nucleotide sequences are deposited in EMBL/Genbank under accession numbers Z97333 (*RHCE*) and Z97334 (*RHD*).

**Figure 4.** Detection of weak D types by PCR-RFLP. Four weak D types harboured point mutations that obliterated restriction sites: Weak D type 1 lacks an *Alw44* site (Panel A), weak D type 3 a *SacI* site (Panel C), weak D type 4 an *AluI* site

(Panel D) and weak D type 6 a *MspI* site (Panel E). In a fifth weak D type, a point mutation introduced a restriction site: Weak D type 2 gained an *AluI* site (Panel B). On the left side of the gels, 100 bp ladders are shown; the position of the 500 bp and 100 bp fragments are indicated on the right side of the panels. For the PCR reaction of panel A, the largest restriction fragment approximation 3,000 bp is not shown.

The example illustrates the invention.

#### **Example: Molecular Analysis of Samples of the Weak D Phenotype**

A method for *RHD* specific sequencing of the ten *RHD* exons and their splice sites was developed (Table 1 and 2). In a sequential analysis strategy, blood samples with weak expression of antigen D were checked by this method, PCR-RFLP (Table 3) and *RHD* PCR-SSP (Gassner et al. 1997). For this purpose, EDTA- or citrate-anticoagulated blood samples were collected from white blood donors and characterized as weak D during donor typing in accordance with published standards („D<sup>u</sup>-test“) (Wissenschaftlicher Beirat der Bundesärztekammer and Bundesgesundheitsamt, 1992) as described (Wagner et al. 1995). D category VI samples were excluded from this study.

**Coding sequence of *RHD* in weak D phenotypes. Sequencing of the ten *RHD* exons from genomic DNA.** DNA was prepared as described previously (Gassner et al. 1997). Nucleotide sequencing was performed with a DNA sequencing unit (Prism dye terminator cycle-sequencing kit with AmpliTaq FS DNA polymerase; ABI 373A, Applied Biosystems, Weiterstadt, Germany). Nucleotide sequencing of genomic DNA stretches representative of all ten *RHD* exons and parts of the promoter (see below) was accomplished using primers (Table 1) and amplification procedures (Table 2) that obviated the need of subcloning steps.

**Control of *RHD* specificity.** *RHD* exons 3 to 7 and 9 carry at least one *RHD* specific nucleotide, which was used to verify the *RHD* origin of the sequences. For exon 1, characteristic nucleotides in the adjacent parts of intron 1 were used (EMBL nucleotide sequence data base accession numbers Z97362 and Z97363). For exon 8, the *RHD* specificity of the PCR amplification was checked by *RHD* non-specific sequencing of the informative exon 9, since exons 8 and 9 were amplified as a single

PCR amplicon (Table 2). Exon 2 and 10 were amplified in an *RHD* specific way (Table 2) based on published *RHD* specific nucleotide sequences used (EMBL nucleotide sequence data base accession numbers U66340 and U66341; Kemp et al. 1996; Le Van Kim et al. 1992); no PCR amplicons were obtained in RhD negative controls. All normal D and weak D samples showed a G at position 654 (Arce et al. 1993) and a C at position 1036 (Le Van Kim et al. 1992), supporting the notion (Cartron, 1996) that the alternatively described C (Le Van Kim et al. 1992) and T (Arce et al. 1993), respectively, were sequencing errors.

**Detection of weak D specific mutations by PCR-RFLP and PCR-RFLP.** PCR-RFLP as well as *RH* PCR-SSP (Gassner et al. 1997) were developed or applied to characterize distinct nucleotide substitutions detected in five *RHD* alleles (see also Tables 3 and 4): The C to G substitution at position 8 led to the loss of a *SacI* restriction site in amplicons obtained with re01 and re11d (G to A at 29, loss of *MspI* site, re01/re11d; C to A at 446, loss of *AluI* site, rb20d/rb21d, T to G at 809, loss of *AluI* site, rf51/re71; G to C at 1154, introduction of *AluI* site, re82/re93). Conditions for the rf51/re71 PCR reaction were as shown in Table 2. The rb20d/rb21d reaction was done with non-proofreading Taq-polymerase (Boehringer Mannheim or Qiagen) with 20 s denaturation at 94°C, 30 s annealing at 60°C and 30 s extension at 72°C. The other PCR reactions were done with non-proofreading Taq-polymerase with 20 s denaturation at 94°C, 30 s annealing at 55°C and 1 min extension at 72°C.

Another four *RHD* alleles were detected by a standard *RH* PCR-SSP<sup>15</sup>: The *RHD*(T201R, F223V) and *RHD*(S182T, K198N, T201R) alleles lacked specific amplicons for *RHD* exon 4, the *RHD*(G307R) and *RHD*(A276P) alleles those for *RHD* exon 6. For all other weak D types, the authenticity of the point mutations was checked by nucleotide sequencing of independent PCR amplicons.

**Sequencing of exons 6 to 9 in *D<sup>IV</sup>* type III.** In *D<sup>IV</sup>* type III exons 6 to 9 were amplified and sequenced using primers that were specific for *RHCE* and *RHD*. Therefore, primer re71 (Table 2) was substituted by primer rb7; primer re621 by rb26; and primer re52 by re74.

Sixteen *RHD* alleles with distinct nucleotide changes coding for amino acid substitutions were identified. (Table 4). One allele represented a typical, yet unpublished, *RHD-CE-D* hybrid allele dubbed hereby *D<sup>IV</sup>* type III. Another allele was

*DHMi* (Liu et al. 1996). Of the remaining 14 alleles, 12 showed single, but distinct previously unknown missense mutations. None of the encoded variant amino acids occurred at the corresponding positions in the RhCE proteins. Two alleles exhibited multiple nucleotide changes typical for the *RHCE* gene, which were interspersed by *RHD* specific sequences.

**Distribution of weak D alleles in whites.** A set of 161 samples with weak expression of antigen D were chosen from random blood donors in South-Western Germany. D category VI samples but no other partial D were excluded by serologic methods. Thus, three samples represented known partial D (*DHMi* (Liu et al. 1996) and D category IV (Lomas et al. 1989)). Without any exception, all samples could be assigned to distinct *RHD* alleles with aberrant coding sequences (Table 5). For the purpose of the present invention, it is proposed that the new molecular weak D types should be referred to by trivial names, e.g. weak D type 1, or by their molecular structures, e.g. *RHD* (V270G). The weak D type 1 was the most frequent known *RHD* allele ( $f=1:277$ ) with aberrant coding sequence, exceeding even the  $D^{VI}$  allele frequency (Wagner et al. 1997).

**Amino acid substitutions in weak D alleles are clustered.** The amino acid substitutions observed in weak D types with single missense mutations were not evenly distributed in the RhD protein (Fig. 1). The majority of substitutions occurred in the region of amino acid positions 267 to 397. Single and multiple amino acid substitutions in smaller portions of the RhD protein around positions 2 to 13, 149, and 179 to 225 (weak D type 4 and 14) were also found in weak D alleles. According to the current RhD loop model, the involved amino acids were positioned in the transmembrane and intracellular protein segments.

**Normal RhD phenotype controls and *RHD* promoter.** Six control samples with normal RhD phenotype showed a normal RhD protein sequence by *RHD* specific sequencing of the ten *RHD* exons. To check for mutations in the *RHD* promoter, a 675 bp region using primer pair rb13 and rb11d were amplified (Table 2). The promoter region was sequenced using primers re02 and re01 starting at nucleotide position -545 relative to the first nucleotide of the start codon. One sample of each weak D type, *DHMi*, and  $D^{IV}$  type III was employed. No deviation from the published *RHD* promoter sequence (Huang, 1996) was found.

**Statistical evidence that missense mutations can cause weak D phenotypes.** The frequency of altered RhD proteins in weak D (158 of 158) and normal D samples (0 of 6) was statistically significantly different ( $p < 0.0001$ , 2x2 contingency table, Fisher's exact test). A normal RhD coding sequence in the weak D phenotype was expected to occur in less than 1.9 % (upper limit of 95% confidence interval, Poisson distribution). It was further excluded that these amino acid substitutions reflected random nucleotide changes only, because of two observations: (i) In the 417 codons of the *RHD* gene, 2,766 missense and 919 silent mutations may occur. If nucleotide changes in weak D alleles were random, silent mutations were expected with a frequency of 0.249. One silent mutation was observed among a total of 18 mutations in weak D alleles ( $p = 0.039$ , binomial distribution). Nonsense mutations were assumed to prevent RhD expression (Avent et al. 1997b) and thus excluded from the calculation. (ii) 1,796 bp of the *RHD* gene were sequenced representing 1,251 bp coding sequence and 545 bp noncoding sequence. If nucleotide changes were random, their occurrence in the noncoding sequence of weak D alleles was expected with a frequency of 0.303. All 18 mutations were, however, located in the coding sequence ( $p = 0.005$ , binomial distribution).

**Haplotype-specific *RHD* polymorphisms.** Introns 3 and 6 were analyzed. To check the *RHD* intron 3 by RFLP, the 3' part of intron 3 using the *RHD* specific primer pair rb46 and rb12 was amplified and the PCR products digested with *Hae*III. To examine TATT tandem repeats in *RHD* intron 6, the full length intron 6 using the *RHD* specific primer pair rf51 and re71 and primer rg62 was amplified used for sequencing.

Polymorphic *RHD* sequences that differed between the prevalent *RHD* alleles of the CDe and cDE haplotypes were detected (Table 6). In *RHD* intron 3, there was a G/C polymorphism that determined a *Hae*III-RFLP at position -371 relative to the intron 3/exon 4 junction. In *RHD* intron 6, there was a variable length TATT tandem repeat starting 1,915 bp 3' of exon 6. In the prevalent *RHD* allele of the CDe haplotype, the *Hae*III restriction site was present and the TATT repeat region comprised 9 repeats. In the prevalent *RHD* allele of the cDE haplotype, the *Hae*III restriction site was absent and the TATT repeat region comprised 8 repeats. Weak D alleles were identical to the prevalent alleles of the same *RH* haplotype in regard to these polymorphisms in intron 3 and 6, with the single exception of weak D type 4 that

showed 13 TATT repeats. It was concluded that weak D alleles evolved independently in the different *RH* haplotypes.

Table 1. Primers used

Name	Nucleotide sequence	Genomic region	Position <sup>1</sup>	Strandedness	RHD specific
ra21	gtgccacttgactgggact	intron 2	2,823 to 2,842	sense	no
rb7	atctctccaagcagaccagcaagc	exon 7	1,022 to 998	antisense	no
rb11	taccttgaattaagcacttcacag	intron 4	161 to 185	sense	yes
rb12	tcctgaacctgctctgtgaagtgc	intron 4	198 to 175	antisense	yes
rb13	ctagagccaaacccacatctcctt	promoter	-675 to -652	sense	no
rb15	ttattggctacttggtgcc	intron 5	-612 to -630	antisense	no
rb20d	tcttggtctccctctct	intron 2	-25 to -8	sense	yes
rb21	aggtccctcctccagcac	intron 3	28 to 11	antisense	no
rb21d	cccagggtccctcctccagcac	intron 3	32 to 11	antisense	no
rb22	gggagatttttcagccag	intron 4	82 to 64	antisense	no
rb24	agaccttggagcaggagt	intron 4	-53 to -34	sense	no
rb25	agcagggaggatgttacag	intron 5	-111 to -93	sense	no
rb26	aggggtgggtagggaatatg	intron 6	-62 to -43	sense	no
rb44	gctgaaatagaagggaatgggagg	intron 7	≈ 3,000	antisense	no
rb46	tggcaagaacctggaccttgactt	intron 3	-1,279 to -1,255	sense	no
rb52	ccagggtgttaagcattgctgtacc	intron 7	≈ - 3,300	sense	yes
re01	atagagaggccagcacaa	promoter	-149 to -132	sense	yes
re02	tgtactatgaggagtcag	promoter	-572 to -554	sense	yes
re11d	agaagatgggggaatcttttct	intron 1	129 to 106	antisense	no
re12d	attagccgggcacggtggca	intron 1	-1,188 to -1,168	sense	yes
re13	actctaatttcataccaccc	intron 1	-72 to -53	sense	no
re23	aaaggatgcaggaggaatgtaggc	intron 2	251 to 227	antisense	no
re31	tgatgaccatcctcaggt	exon 3	472 to 455	antisense	yes
re617	tctcagctcactgcaacctc	intron 6	1,998 to 2,017	sense	no
re621	catcccccttggtgcc	intron 6	- 102 to -85	sense	yes
re71	accagcaagctgaagtgtagcc	exon 7	1,008 to 985	antisense	yes
re73	ctttttgtccctgatgacc	intron 7	- 67 to -48	sense	no
re74	tatccatgaggtgctgggaac	intron 7	≈ -200	sense	no
re75	aaggtaggggtggacag	intron 7	≈ 120	antisense	yes
re82	aaaaatcctgtgtccaaac	intron 8	≈ -45	sense	yes
re83	gagattaaaaatcctgtgtcca	intron 8	≈ -50	sense	no
re91	caagagatcaagccaaaatcagt	intron 9	≈ -40	sense	no
re93	caccgcgatgtcagactattggc	intron 9	≈ 300	antisense	no
rf51	caaaaacccattctccc	intron 5	-332 to -314	sense	no
rg62	tgtattccaggcagaaggc	intron 6	1,736 to 1,755	sense	no
rh5	gcacagagacggacacag	5' UTR <sup>2</sup>	-19 to -2	sense	no
rh7	acgtacaaatgcaggcaac	3' UTR <sup>2</sup>	1,330 to 1,313	antisense	no
rr1	tgttgagagaggggtgatg	5' UTR	-60 to -41	sense	no
rr3	cagtcgtgtgttaccagatg	3' UTR	1,512 to 1,492	antisense	yes
rr4	agcttactggatgaccacca	3' UTR	1,541 to 1,522	antisense	yes

<sup>1</sup> The positions of the synthetic oligonucleotides are indicated relative to their distances from the first nucleotide position of the start codon ATG for all primers in the promoter and in the exons including the 3' untranslated part of exon 10, or relative to their adjacent exon/intron boundaries for all other primers. Primer ra21 was reported previously (Poulter et al. 1996).

<sup>2</sup> 5' UTR: 5' untranslated region of exon 1; UTR: 3' untranslated region of exon 10.

**Table 2.** Sequencing method for all ten *RHD* exons from genomic DNA

<i>RHD</i> exon	PCR primers		<i>RHD</i>	PCR conditions <sup>1</sup>		Sequencing	<i>RHD</i>
	Sense	Antisense	specific <sup>2</sup>	Extension	Annealing	primers	specific <sup>2</sup>
Exon 1	rb13	rb11d	no	10 min	60°C	re01	yes
Exon 2	re12d	re23	yes	3 min	65°C	re13	no
Exon 3	ra21	rb21	no	10 min	60°C	re31 and rb20d	yes
Exon 4	rb46	rb12	yes	10 min	60°C	rb22	no
Exon 5	rb11	rh2	yes	10 min	60°C	rb24	no
Exon 6	rf51	re71	yes	10 min	60°C	rb25	no
Exon 7	re617	rb44	no	10 min	60°C	re621 and re75	yes
Exon 8	rb52	rb93	yes	10 min	60°C	re73	no
Exon 9	rb52	rb93	yes	10 min	60°C	re82/re83	yes/no
Exon 10	re91	rr4	yes	10 min	60°C	rr3/ rh7	yes/no

<sup>1</sup> Primers were used at a concentration of 1 nM in the Expand High Fidelity PCR System (Boehringer Mannheim, Mannheim, Germany). In the exon 10 PCR, the concentration of MgCl<sub>2</sub> was 2.0 nM. Denaturation was 20 s at 92°C, annealing 30 s, elongation temperature 68°C. Elongation time was increased by 20 s for each cycle after the 10th cycle, except for the re12d/re23 primer pair.

<sup>2</sup> To achieve *RHD* specificity for genomic nucleotide sequencing, the PCR primer pairs or the sequencing primer or both must not concurrently detect *RHCE*-derived nucleotide sequences. Primer sequences are given in Table 1.



Table 3. PCR-RFLP analysis of five *RHD* alleles

Allele	Substitution	PCR primers		Restriction enzyme
<i>RHD</i> (S3C)	8 C→G	re01	re11d	SacI
<i>RHD</i> (R10Q)	29 G→A	re01	re11d	MspI
<i>RHD</i> (A149D)	446 C→A	rb20d	rb21	AluI
<i>RHD</i> (V270G)	809 T→G	rf51	re71	Alw44I
<i>RHD</i> (G385A)	1154 G→C	re82	re93	AluI

<sup>1</sup> Conditions for the rf51/re71 PCR reaction as shown in Table 2. All other PCR reactions were done with non-proofreading Taq-polymerase (Boehringer Mannheim) with 20 s denaturation at 94°C, 30 s annealing at 55°C and 1 min extension at 72°C. Examples for these PCR-RFLPs of weak D types 1, 2, 3, 5, 6, are shown in Fig. 4.

Table 4. Molecular basis of weak RhD phenotypes

Allele	Effect on Nucleotide change	Exons protein sequence	Predicted localization involved in	the cell
<i>RHD</i> (S3C)	C→G at 8	Ser to Cys at 3	1	IC
<i>RHD</i> (R10Q)	G→A at 29	Arg to Gln at 10	1	IC
<i>RHD</i> (W16C, T201R, F223V)	G→C at 48,	Trp to Cys at 16	1	TM
	C→G at 602,	Thr to Arg at 201	4	IC
	T→G at 667,	Phe to Val at 223	5	TM
	G→A at 819	no change	6	-
	C→T at 340	Arg to Trp at 114	3	TM
<i>RHD</i> (R114W)	C→A at 446	Ala to Asp at 149	3	TM
<i>RHD</i> (A149D)	T→A at 544,	Ser to Thr at 182	4	TM
<i>RHD</i> (S182T, K198N, T201R)	A→T at 594,	Lys to Asn at 198	4	IC
	C→G at 602	Thr to Arg at 201	4	IC
	C→G at 602,	Thr to Arg at 201	4	IC
	T→G at 667,	Phe to Val at 223	5	TM
	G→A at 819	no change	6	-
<i>RHD</i> (T201R, F223V)	T→C at 858	Trp to Cys at 220	5	TM
	T→G at 809	Val to Gly at 270	6	TM
	G→C at 826	Ala to Pro at 276	6	TM
	G→A at 830	Gly to Glu at 277	6	TM
	G→A at 845	Gly to Asp at 282	6	TM
<i>RHD</i> (G282D)	G→C at 880	Ala to Pro at 294	6	TM
<i>RHD</i> (A294P)	G→T at 885	Met to Ile at 295	6	TM
<i>RHD</i> (M295I)	G→A at 919	Gly to Arg at 307	6	IC
<i>RHD</i> (G307R)	G→A at 1016	Gly to Glu at 339	7	TM
<i>RHD</i> (G339E)	G→C at 1154	Gly to Ala at 385	9	TM
<i>RHD</i> (G385A)	T→C at 1177	Trp to Arg at 393	9	IC
<i>RHD</i> (W393R)				
<i>DHMi</i>	C→T at 848	Thr to Ile at 283	6	EF
<i>D<sup>vi</sup></i> type III	<i>RHD</i> -CE(6-9)-D	multiple	6 to 9	EF/TM/IC

IC - intracellular, TM - transmembranous, EF - exofacial

Table 5. Proposed nomenclature for *RHD* alleles coding for weak D phenotypes and their minimal population frequencies

Trivial name	Molecular basis (allele)	n <sup>1</sup>	Phenotype frequency <sup>2</sup>	Minimal population frequency <sup>3</sup>	
				phenotype	haplotype
weak D type 1	<i>RHD</i> (V270G)	95	70.29%	0.2964%	0.003606 (1:277)
weak D type 2	<i>RHD</i> (G385A)	43	18.01%	0.0759%	0.000924 (1:1,082)
weak D type 3	<i>RHD</i> (S3C)	7	5.19%	0.0219%	0.000266 (1:3,759)
weak D type 4 <sup>*</sup>	<i>RHD</i> (T201R,F223V)	6	1.30%	0.0055%	0.000067 (1:14,925)
weak D type 5	<i>RHD</i> (G307R)	1	0.74%	0.0031%	0.000038 (1:26,316)
weak D type 6	<i>RHD</i> (R10Q)	1	0.74%	0.0031%	0.000038 (1:26,316)
weak D type 7	<i>RHD</i> (G339E)	1	0.74%	0.0031%	0.000038 (1:26,316)
weak D type 8	<i>RHD</i> (A294P)	1	0.42%	0.0017%	0.000021 (1:47,619)
weak D type 9	<i>RHD</i> (A149D)	1	0.42%	0.0017%	0.000021 (1:47,619)
weak D type 10	<i>RHD</i> (W393R)	1	0.42%	0.0018%	0.000021 (1:47,619)
weak D type 11	<i>RHD</i> (M295I)	1	0.22%	0.0009%	0.000011 (1:90,909)
weak D type 12	<i>RHD</i> (G277E)	0	-	-	-
weak D type 13	<i>RHD</i> (A276P)	0	-	-	-
weak D type 14	<i>RHD</i> (S182T,K198N,T201R)	0	-	-	-
weak D type 15	<i>RHD</i> (G282D)	0	-	-	-
weak D type 16	<i>RHD</i> (W220R)	0	-	-	-
weak D type 17	<i>RHD</i> (R114W)	0	-	-	-
DHMi	<i>RHD</i> (T283I)	2	0.84%	0.0035%	0.000043 (1:23,256)
D <sup>iv</sup> type III	<i>RHD</i> -CE(6-9)-D	1	0.60%	0.0025%	0.000031 (1:32,258)
Total		161	100%	0.4185%	0.005094

- weak D type 4 can be subdivided in two forms:
- weak D type 4a; see weak D type 4, in the Table
- weak D type 4b *RHD*(W16C, T201R, F223V)

Continued on next page

Table 5 - Continued

1	Number of samples observed among 161 blood samples with weak antigen D expression. Types 12 to 17 were not detected among these blood samples, but found independently.
2	The phenotype frequencies among weak D samples were calculated adjusting for the frequencies of the serologic weak D phenotypes (Wagner et al 1995). cDE weak D samples were assumed to be cDE/cDE.
3	Phenotype frequencies in the population were calculated from the population frequency of the weak D phenotype in South-Western Germany (Wagner et al 1995). These are minimal estimates, because some samples with only moderately weakened D expression may have been grouped to normal strength D. Haplotype frequencies were calculated using a haplotype frequency of 0.411 for RhD negative haplotypes (Wagner et al 1995) assuming that all weak D samples were heterozygous.

Table 6. *RHD* polymorphisms in *RHD* genes of various haplotypes

Haplotype <sup>1</sup> Allele	HaeIII site in intron 3	TATT-repeat in intron 6
CDe prevalent <i>RHD</i>	present	9
CDe weak <i>D</i> type 1	present	9
CDe weak <i>D</i> type 3	present	9
CDe weak <i>D</i> type 5	present	9
CDe weak <i>D</i> type 6	present	9
CDe weak <i>D</i> type 7	present	9
CDe weak <i>D</i> type 12	present	9
CDe weak <i>D</i> type 13	present	9
CDe weak <i>D</i> type 17	N.D.	9
CDe <i>D<sup>v</sup></i> type III	present	<sup>2</sup>
cDE prevalent <i>RHD</i> <sup>3</sup>	absent	8
cDE weak <i>D</i> type 2	absent	8
cDE weak <i>D</i> type 8	absent	8
cDE weak <i>D</i> type 9	absent	8
cDE weak <i>D</i> type 10	absent	8
cDE weak <i>D</i> type 14	absent	8
cDE weak <i>D</i> type 15	absent	8
cDE weak <i>D</i> type 16	absent	8
cDE <i>DHMi</i>	absent	8
c(W16C)Devalent <i>RHD</i>	present	9
c(W16C)De weak <i>D</i> type 4	present	13
cDe prevalent <i>RHD</i> <sup>4</sup>	mostly present	8 or 9
cDe weak <i>D</i> type 11	present	8

<sup>1</sup> The haplotype association of the HaeIII site was tested in 10 CCDee, 8 ccDEE, 10 cc(W16C)De and 10 ccDee samples. The association of the TATT repeat was tested in 3 CCDee, 3 ccDEE, 1ccDEe, 2 cc(W16C)De and 2 ccDee samples.

<sup>2</sup> Intron 6 derived from *RHCE* due to a gene conversion.

Continued on next page

Table 6 - Continued

3	Six of seven alleles investigated showed 8 repeats, one 9 repeats.
4	The HaeIII site was present in 8 of 10 samples tested. Samples with HaeIII site showed 9 TAT repeats, samples without HaeIII site 8 repeats
	N D = Not determined

**Table 7.** Predicted localization of RhD protein segments relative to the red blood cells' membrane<sup>1</sup>

Range of amino acids	Intra-cellular	Trans-membranous	Exofacial	Length (amino acids)
1 - 11	X			10 <sup>2</sup>
12 - 31		X		20
32 - 53			X	22
54 - 71		X		18
72 - 75	X			4
76 - 93		X		18
94 - 110			X	17
111 - 130		X		20
131 - 134	X			4
135 - 153		X		19
154 - 169			X	16
170 - 187		X		18
188 - 207	X			20
208 - 225		X		18
226 - 238			X	13
239 - 256		X		18
257 - 264	X			8
265 - 282		X		18
283 - 286			X	4
287 - 306		X		20
307 - 333	X			27
334 - 351		X		18
352 - 370			X	19
371 - 388		X		18
389 - 417	X			29
Total	5 loops and 2 segments	12 helices	6 loops	416 <sup>2</sup>

<sup>1</sup> Localization of the amino- and carboxyterminal protein end according to Avent et al. {*J. Biol. Chem.* 1992} and Hermand et al. {*Blood* 1993}. The transmembranous helices were predicted by PHDhtm {[www.embl-heidelberg.de/predictprotein/predictprotein.html](http://www.embl-heidelberg.de/predictprotein/predictprotein.html)}, the helix at positions 371 to 388 by TMPred {[ulrec3.unil.ch/software/TMPRED\\_form.html](http://ulrec3.unil.ch/software/TMPRED_form.html)}.

<sup>2</sup> The amino acid (methionine) at position 1 is not expressed in the mature RhD protein as shown by amino acid sequencing {Avent et al. *Biochem. J.* 1988}.

**Table 8.** Sample RhD epitope densities for weak D types.

weak D	RhD epitope density (RhD antigens/red cell)
type 3	1,500
type 1	900
type 2	500
type 12	<100

One sample of each weak D type was tested with a polyclonal anti-D (Lorne Laboratories Ltd., Redding, Berkshire, England) as described previously (Flegel and Wagner 1996). Similar results were obtained by monoclonal anti-D (BS228, Biotest AG, Dreieich, Germany; and P3x290, Diagast, Lille, France).



## References

- Agre, P.C., Davies, D.M., Issitt, P.D., Lamy, B.M., Schmidt, P.J., Treacy, M., and Vengelen-Tyler, V. 1992. A proposal to standardize terminology for weak D antigen [Letter]. *Transfusion* 32:86-87.
- Arce, M.A., Thompson, E.S., Wagner, S., Coyne, K.E., Ferdman, B.A., and Lublin, D.M. 1993. Molecular cloning of RhD cDNA derived from a gene present in RhD-positive, but not RhD-negative individuals. *Blood* 82:651-655.
- Aubin, J.T., Le Van Kim, C., Mouro, I., Colin, Y., Bignozzi, C., Brossard, Y., and Cartron, J.P. 1997. Specificity and sensitivity of RHD genotyping methods by PCR-based DNA amplification. *British Journal of Haematology* 98:356-364.
- Avent, N.D., Butcher, S.K., Liu, W., Mawby, W.J., Mallinson, G., Parsons, S.F., Anstee, D.J., Tanner, M.J. 1992. Localization of the C termini of the Rh(rhesus) polypeptides to the cytoplasmic face of the human erythrocyte membrane. *J. Biol. Chem.* 267:15134-15139.
- Avent, N.D., Jones, J.W., Liu, W., Scott, M.L., Voak, D., Flegel, W.A., Wagner, F.F., and Green, C. 1997a. Molecular basis of the D variant phenotypes DNU and DII allows localization of critical amino acids required for expression of Rh D epitopes epD3, 4 and 9 to the sixth external domain of the Rh D protein. *British Journal of Haematology* 97:366-371.
- Avent, N.D., Martin, P.G., Armstrong-Fisher, S.S., Liu, W., Finning, K.M., Maddocks, D., and Urbaniak, S.J. 1997b. Evidence of genetic diversity underlying Rh D negative, weak D (D<sup>u</sup>) and partial D phenotypes as determined by multiplex PCR analysis of the RHD gene. *Blood* 89:2568-2577.
- Avent, N.D., Rigwell, K., Mawby, W.J., Tanner, M.J., Anstee, D.J., Kumpel, B. 1988. Protein-sequence studies on Rh-related polypeptides suggest the presence of at least two groups of proteins which associate in the human red-cell membrane. *Biochem. J.* 256:1043-1046.
- Beckers, E.A., Faas, B.H., Ligthart, P., Overbeeke, M.A., von dem Borne, A.E., van der Schoot, C.E., and van Rhenen, D.J. 1997. Lower antigen site density and weak D immunogenicity cannot be explained by structural genomic abnormalities or regulatory defects of the RHD gene. *Transfusion* 37:616-623.
- Beckers, E.A.M., Faas, B.H.W., Overbeeke, M.A.M., von dem Borne, A.E.G.K., van Rhenen, D.J., and van der Schoot, C.E. 1995. Molecular aspects of the weak-D phenotype. *Transfusion* 35:50S.
- Bowman, J.M. 1998. RhD hemolytic disease of the newborn. *N. Engl. J. Med.* 339:1775-1777.

- Cartron, J.-P. 1996. Rh DNA - coordinator's report. *Transfusion Clinique et Biologique* 3:491-495.
- Ceppellini, R., Dunn, L.C., and Turry, M. 1955. An interaction between alleles at the Rh locus in man which weakens the reactivity of the Rho factor (D<sup>o</sup>). *Proceedings of the National Academy of Sciences U.S.A.* 41:283-288.
- Cherif-Zahar, B., Raynal, V., Gane, P., Mattei, M.G., Bailly, P., Gibbs, B., Colin, Y., and Cartron, J.-P. 1996. Candidate gene acting as a suppressor of the RH locus in most cases of Rh-deficiency. *Nature Genetics* 12:168-173.
- Flegel, W. A., Müller, T.H., Schunter, F., Gassner, C., Schönitzer, D., and Wagner, F.F. 1997. D category VI type III: A D-Ce(3-6)-D hybrid protein with normal RhD antigen density on red cells [Abstract]. *Transfusion* 37S:101S
- Flegel, W.A. and Wagner, F.F. 1996. RHD epitope density profiles of RHD variant red cells analyzed by flow cytometry. *Transfusion Clinique et Biologique* 3:429-431.
- Fukumori, Y., Hori, Y., Ohnoki, S., Nagao, N., Shibata, H., Okubo, Y., and Yamaguchi, H. 1997. Further analysis of Del (D-elute) using polymerase chain reaction (PCR) with RHD gene-specific primers. *Transfusion Medicine* 7:227-231.
- Gassner, C., Schmarda, A., Kilga-Nogler, S., Jenny-Feldkircher, B., Rainer, E., Müller, T.H., Wagner, F.F., Flegel, W.A., and Schönitzer, D. 1997. RhesusD/CE typing by polymerase chain reaction using sequence-specific primers. *Transfusion* 37:1020-1026.
- Hasekura, H., Ota, M., Ito, S., Hasegawa, Y., Ichinose, A., Fukushima, H., and Ogata, H. 1990. Flow cytometric studies of the D antigen of various Rh phenotypes with particular reference to D<sup>o</sup> and D<sup>el</sup>. *Transfusion* 30:236-238.
- Hernand, P., Mouro, I., Huet, M., Bloy, C., Suyama, K., Goldstein, J., Cartron, J.P., Bailly, P. 1993. Immunochemical characterization of rhesus proteins with antibodies raised against synthetic peptides. *Blood* 82:669-676.
- Huang, C.H. 1996. Alteration of RH gene structure and expression in human dCCee and DCW- red blood cells: phenotypic homozygosity versus genotypic heterozygosity. *Blood* 88:2326-2333.
- Huang, C.H. 1997. Molecular insights into the Rh protein family and associated antigens. *Current Opinion in Hematology* 4:94-103.
- Huang, C.H., Chen, Y., and Reid, M. 1997. Human D(IIIa) erythrocytes: RhD protein is associated with multiple dispersed amino acid variations. *Am. J Hematol.* 55:139-145.

- Issitt, P.D. and Telen, M.J. 1996. D, weak D (D<sup>w</sup>), and partial D: the molecular story unfolds. *Transfusion* 36:97-100.
- Jones, J.W., Finning, K.M., Mattock, R., Voak, D., Scott, M.L., and Avent, N.D. 1997. The serological profile and molecular basis of a new partial D phenotype, DHR. *Vox Sanguinis* 73:252-256.
- Jones, J.W., Lloyd-Evans, P., and Kumpel, B.M. 1996. Quantitation of Rh D antigen sites on weak D and D variant red cells by flow cytometry. *Vox Sanguinis* 71:176-183.
- Kajii, E., Umenishi, F., Omi, T., and Ikemoto, S. 1995. Intricate combinatorial patterns of exon splicing generate multiple Rh- related isoforms in human erythroid cells. *Human Genetics* 95:657-665.
- Kemp, T.J., Poulter, M., and Carritt, B. 1996. A recombination hot spot in the Rh genes revealed by analysis of unrelated donors with the rare D<sup>--</sup> phenotype. *American Journal of Human Genetics* 59:1066-1073.
- Le Van Kim, C., Mouro, I., Cherif-Zahar, B., Raynal, V., Cherrier, C., Cartron, J.P., and Colin, Y. 1992. Molecular cloning and primary structure of the human blood group RhD polypeptide. *Proceedings of the National Academy of Sciences U.S.A.* 89:10925-10929.
- Leader, K.A., Kumpel, B.M., Poole G.D., Kirkwood, J.T., Merry A.H. and Bradley, B.A. 1990. Human monoclonal anti-D with reactivity against category DVI cells used in blood grouping and determination of the incidence of the category DVI phenotype in the DU population. *Vox Sang* 58:(2):106-11
- Legler, T.J., Blaschke, V., Bustami, N., Malekan, M., Schwartz, D.W.M., Mayr, W.R., Panzer, S., and Köhler, M. 1997. RHD genotyping on exons 2, 5, 7, intron 4 and the 3' non-coding region in D<sup>weak</sup>, D<sup>VI</sup>, DFR and D<sup>--</sup> individuals. *Transfusion* 37:100S
- Liu, W., Jones, J.W., Scott, M.L., Voak, D., and Avent, N.D. 1996. Molecular analysis of two D-variants, D<sup>HMi</sup> and D<sup>HMi</sup> [Abstract]. *Transfusion Medicine* 6(suppl 2):21
- Lomas, C., Grässmann, W., Ford, D., Watt, J., Gooch, A., Jones, J., Beolet, M., Stern, D., Wallace, M., and Tippett, P. 1994. FPTT is a low-incidence Rh antigen associated with a "new" partial Rh D phenotype, DFR. *Transfusion* 34:612-616.
- Lomas, C., McColl, K., and Tippett, P. 1993. Further complexities of the Rh antigen D disclosed by testing category D<sup>w</sup> cells with monoclonal anti-D. *Transfusion Medicine* 3:67-69.
- Lomas, C., Tippett, P., Thompson, K.M., Melamed, M.D., and Hughes-Jones, N.C. 1989. Demonstration of seven epitopes on the Rh antigen D using human monoclonal anti-D antibodies and red cells from D categories. *Vox Sanguinis* 57:261-264.

- Mollison, P. L., Engelfriet, C.P., and Contreras, M. 1993. *Blood transfusion in clinical medicine*. 9th ed. London: Blackwell Scientific Publications.
- Moore, B.P.L. 1984. Does knowledge of D<sup>u</sup> status serve a useful purpose? *Vox Sanguinis* 46S1:95-97.
- Mourant, A. E., Kopec, A.C., and Domaniewska-Sobczak, K. 1976. *The distribution of the human blood groups and other polymorphisms*. 2nd ed. London: Oxford University Press.
- Nelson, M., Barrow, L.A., Popp, H., and Gibson, J. 1995. Some observations on D antigen expression of D-positive and 'weak D- positive' red cells as assessed by flow cytometry. *Vox Sanguinis* 69:152-154.
- Nicholson, G., Lawrence, A., Ala, F.A., and Bird, G.W.G. 1991. Semi-quantitative assay of D antigen site density by flow cytometric analysis. *Transfusion Medicine* 1:87-90.
- Poulter, M., Kemp, T.J., and Carritt, B. 1996. DNA-based Rhesus typing: simultaneous determination of RHC and RHD status using the polymerase chain reaction. *Vox Sanguinis* 7:164-168.
- Roubinet, F., Apoil, P.A., and Blancher, A. 1996. Frequency of partial D phenotypes in the south western region of France. *Transfusion Clinique et Biologique* 3:247-255.
- Rouillac, C., Gane, P., Cartron, J.-P., Le Pennec, P.Y., and Colin, Y. 1996. Molecular basis of the altered antigenic expression of RhD in weak D (D<sup>u</sup>) and RhC/e in R<sup>N</sup> phenotypes. *Blood* 87:4853-4861.
- Rouillac, C., Le Van Kim, C., Beolet, M., Cartron, J.P., and Colin, Y. 1995. Leu110Pro substitution in the RhD polypeptide is responsible for the D<sup>VI</sup> category blood group phenotype. *American Journal of Hematology* 49:87-88.
- Salmon, C., Cartron, J.-P., and Rouger, P. 1984. *The human blood groups*. New York: Masson.
- Scott, M. 1996. Rh serology - coordinator's report. *Transfusion Clinique et Biologique* 3:333-337.
- Siegel, D.L., Silberstein, L.E. 1994. Expression and characterization of recombinant anti-Rh(D) antibodies on filamentous phage: a model system for isolating human red blood cell antibodies by repertoire cloning. *Blood* Apr. 15:83(8):2334-44
- Stratton, F. 1946. A new Rh allelomorph. *Nature* 158:25

- Tazzari, P.L., Bontadini, A., Belletti, D., Malferrari, F., and Conte, R. 1994. Flow cytometry: a tool in immunohematology for D+<sup>w</sup> (D<sup>w</sup>) antigen evaluation? *Vox Sanguinis* 67:382-386.
- Tippett, P. and Sanger, R. 1962. Observations on subdivisions of Rh antigen D. *Vox Sanguinis* 7:9-13.
- Tippett, P. and Sanger, R. 1977. Further observations on subdivisions of the Rh antigen D. *Das Ärztliche Laboratorium* 23:476-480.
- Wagner, F.F. 1994. Influence of Rh phenotype on the antigen density of C, c, and D: flow cytometric study using a frozen standard red cell. *Transfusion* 34:671-676.
- Wagner, F.F., Hillesheim, B., and Flegel, W.A. 1997. D-Kategorie VII beruht einheitlich auf der Aminosäuresubstitution Leu(110)Pro. *Beiträge zur Infusionstherapie und Transfusionsmedizin* 34:220-223.
- Wagner, F.F., Kasulke, D., Kerowgan, M., and Flegel, W.A. 1995. Frequencies of the blood groups ABO, Rhesus, D category VI, Kell, and of clinically relevant high-frequency antigens in South-Western Germany. *Infusionstherapie und Transfusionsmedizin* 22:285-290.
- Wissenschaftlicher Beirat der Bundesärztekammer and Bundesgesundheitsamt. 1992. *Richtlinien zur Blutgruppenbestimmung und Bluttransfusion*. Köln: Deutscher Ärzte-Verlag.

## CLAIMS

1. A nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, in its transmembrane and/or intracellular regions.
2. A nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule
  - (a) carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, in amino acid positions 2-16, 114-149, 179-225 or/and 267 to 397 with the proviso that said D antigen does carry not a single missense mutation leading to a substitution of phenylalanine in amino acid position 223 by valine or of threonine in position 283 by isoleucine; or
  - (b) carrying a gene conversion involving exons 6 to 9 which are replaced by the corresponding exons of the *RHCE* gene.
3. The nucleic acid molecule of claim 1 or 2 wherein said missense mutation causes an amino acid substitution in position 3, 10, 16, 114, 149, 182, 198, 201, 220, 223, 270, 276, 277, 282, 294, 295, 307, 339, 385 and 393 or a combination of/or involving said substitutions.
4. The nucleic acid molecule of claim 3 wherein said amino acid substitution in position 3 is from Ser to Cys, in position 10 from Arg to Gln, in position 16 from Trp to Cys, in position 114 from Arg to Trp, in position 149 from Ala to Asp, in position 182 from Ser to Thr, in position 198 from Lys to Asn, in position 201 from Thr to Arg, in position 220 from Trp to Arg, in position 223 from Phe to Val, in position 270 from Val to Gly, in position 276 from Ala to Pro, in position 277 from Gly to Glu, in position 282 from Gly to Asp, in position 294 from Ala to Pro, in position 295 from Met to Ile, in position 307 from Gly to Arg, in position 339 from Gly to Glu, in position 385 from Gly to Ala and in position 393 from Trp to Arg.

5. The nucleic acid molecule of any one of claims 1 to 4 wherein said missense mutation occurs in nucleotide position 8, 29, 48, 340, 446, 544, 594, 602, 658, 667, 809, 819, 826, 830, 845, 880, 885, 919, 1016, 1154 or 1177 or in a combination of said positions.
6. The nucleic acid molecule of claim 5 wherein said missense mutation in position 8 is from C to G, in position 29 from G to A, in position 48 from G to C, in position 340 from C to T, in position 446 from C to A, in position 544 from T to A, in position 594 from A to T, in position 602 from C to G, in position 658 from T to C, in position 667 from T to G, in position 809 from T to G, in position 819 from G to A, in position 826 from G to C, in position 830 from G to A, in position 845 from G to A, in position 880 from G to C, in position 885 from G to T, in position 919 from G to A, in position 1016 from G to A, in position 1154 from G to C and in position 1177 from T to C.
7. The nucleic acid molecule of claim 3 or 4 wherein said combination of substitutions is in positions 182, 198 and 201 and is preferably S182T, K198N, T201R or in position 201 and 223 and is preferably T201R and F223V, or in position 16, 201 and 223 and is preferably W16C, T201R and F223V.
8. The nucleic acid molecule of claim 5 or 6 wherein said combination of missense mutations comprises positions 544, 594 and 602 and is preferably T→A at position 544, A→T at position 594 and C→G at position 602 or comprises positions 602, 677 and 819 and is preferably C→G at position 602, T→G at position 667 and G→A at position 819 or comprises positions 48, 602, 667 and 819 and is preferably G→C at position 48, C→G at position 602, T→G at position 667 and G→A at position 819.
9. The nucleic acid molecule of any one of claims 1 to 8 which is mRNA or genomic DNA.
10. A vector comprising the nucleic acid molecule of any one of claims 1 to 9.

11. A host transformed with the vector of claim 10.
12. A method of producing a Rhesus D antigen contributing to the weak D phenotype comprising culturing the host of claim 11 under suitable conditions and isolating the Rhesus D antigen produced.
13. A Rhesus D antigen encoded by the nucleic acid molecule of any one of claims 1 to 9 or produced by the method of claim 12.
14. An oligonucleotide hybridizing under stringent conditions to a portion of the nucleic acid molecule of any one of claims 1 to 9 comprising said at least one missense mutation or to the complementary portion thereof or hybridizing to a breakpoint of the gene conversion identified in claim 2.
15. An antibody or aptamer or phage specifically binding to the Rhesus D antigen of claim 13.
16. An antibody or aptamer or phage specifically binding to the wild type Rhesus D antigen or to aberrant Rhesus D antigens but not to the Rhesus D antigen of claim 13.
17. A method for testing for the presence of a nucleic acid molecule encoding a Rhesus D antigen contributing to the weak D phenotype in a sample comprising hybridizing the oligonucleotide of claim 14 or an oligonucleotide hybridizing to a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T



under stringent conditions to nucleic acid molecules comprised in the sample obtained from a human and detecting said hybridization.

18. The method of claim 17 further comprising digesting the product of said hybridization with a restriction endonuclease and analyzing the product of said digestion.
19. A method of testing for the presence of a nucleic acid encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising determining the nucleic acid sequence of at least a portion of the nucleic acid molecule of any one of claims 1 to 9, said portion encoding at least one of said missense mutations or a breakpoint of said gene conversion or a nucleic acid molecule encoding a Rhesus D antigen contributing to the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T.
20. The method of claim 19 further comprising, prior to determining said nucleic acid sequence, amplification of at least said portion of said nucleic acid molecule.
21. A method for testing for the presence of a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising carrying out an amplification reaction wherein at least one of the primers employed in said amplification reaction is the oligonucleotide of claim 14 or an oligonucleotide hybridizing to a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid

substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is form T to G and in position 848 from C to T and assaying for an amplification product.

22. The method of claim 20 or 21 wherein said amplification or amplification reaction is or is effected by the polymerase chain reaction (PCR).
23. A method for testing for the presence of a Rhesus D antigen contributing to or indicative of the weak D phenotype in a sample comprising assaying a sample obtained from a human for specific binding to the antibody or aptamer or phage of claim 15 or to an antibody or aptamer or phage to a Rhesus D antigen contributing to or indicative of the weak D phenotype and encoded by a nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is form T to G and in position 848 from C to T.
24. A method of testing a sample for the presence of wild type Rhesus D antigen and the absence of the Rhesus D antigen of claim 13 comprising assaying a sample obtained from a human for specific binding to the antibody or aptamer or phage of claim 16.
25. The method of any one of claims 17 to 24 wherein said sample is blood, serum, plasma, fetal tissue, saliva, urine, mucosal tissue, mucus, vaginal tissue, fetal tissue obtained from the vagina, skin, hair, hair follicle or another human tissue.
26. The method of 25 comprising enrichment of fetal cells or extraction of fetal DNA or mRNA from material tissue, like peripheral blood, serum or plasma.

27. The method of any one of claims 17 to 26 wherein said nucleic acid molecule or proteinaceous material from said sample is fixed to a solid support.
28. The method of claim 27 wherein said solid support is a chip.
29. Use of the nucleic acid molecule of any one of claims 1 to 9 or of a nucleic acid molecule encoding a Rhesus D antigen contributing to or indicative of the weak D phenotype, said nucleic acid molecule carrying at least one missense mutation, as compared to the wild type Rhesus D antigen, said missense mutation causing an amino acid substitution in position 223 or 283 which is in position 223 preferably from Phe to Val and in position 283 preferably from Thr to Ile, said missense mutation further preferably occurring in nucleotide position 667 or 848 wherein most preferably said mutation in position 667 is from T to G and in position 848 from C to T or of a combination thereof for the analysis of a weak Rhesus D phenotype.
30. Use of the nucleic acid molecule of any one of claims 1 to 9, the vector of claim 10 or the Rhesus D antigen of claim 13 for the assessment of the affinity, avidity and/or reactivity of monoclonal antibodies or of polyclonal antisera preferably anti-D antisera, anti-globulin or anti-human-globulin antisera.
31. A method for the characterization of the monoclonal antibodies or polyclonal antisera or of a preparation thereof said method comprising
  - (a) testing the nucleic acid of sample of a proband for the presence of a mutation as defined in any one of claims 1 to 24;
  - (b) correlating, on the basis of the mutation status and the allelic status of the *RHD* gene, the nucleic acid with the RhD antigen density on the surface of red blood cells of said proband;
  - (c) reacting said monoclonal antibodies or polyclonal antisera or said preparation thereof with a cell carrying the RhD antigen on its surface;
  - (d) characterizing said monoclonal antibodies or polyclonal antisera or said preparation thereof on the basis of the results obtained in step (c).

32. The method of claim 31 wherein said characterization comprises the determination of reactivity, sensitivity, avidity, affinity, specificity and/or other characteristics of antibodies and antisera.
33. The method of claim 31 or 32 wherein said cell carrying the RhD antigen on its surface is a red blood cell.
34. Use of an aptamer, phage, monoclonal antibody or a polyclonal antisera or a preparation thereof as characterized in claim 15 or 16 for RhD antigen determination.
35. Use of claim 34 wherein said RhD antigen determination is effected in connection with blood group typing.
36. A preparation comprising the antibody or aptamer or phage of claim 15 or 16.
37. A method of identifying an antibody  $V_H$  or  $V_L$  chain or a combination thereof or an aptamer specifically binding to a weak D polypeptide of claim 13 comprising
  - (a) contacting the weak D polypeptide of claim 13 with a phage library displaying  $V_H$  or  $V_L$  chains or combinations thereof on the surface of the phage or with aptamers;
  - (b) identifying phage or aptamers that bind to said weak D polypeptide; and optionally
  - (c) repeating steps (a) and (b) one or more times.
38. A method of identifying a monoclonal antibody specifically binding to a weak D polypeptide/antigen of claim 13 comprising
  - (a) contacting the weak D polypeptide of claim 13 with one or more monoclonal antibodies;
  - (b) identifying monoclonal antibodies that bind to said weak D polypeptide; and optionally
  - (c) repeating steps (a) and (b) one or more times.


39. A method of identifying an antibody  $V_H$  or  $V_L$  chain or a combination thereof or an aptamer specifically binding to a weak D polypeptide/antigen of claim 13 comprising
- (a) contacting the weak D polypeptide and
  - (aa) a second or more weak D polypeptide(s) and/or
  - (ab) a normal D polypeptide
- wherein the second or more weak D polypeptide(s) and/or the normal D polypeptide are present in a molar mass that is higher, equal or less than the weak D polypeptide of (a) with a phage library displaying  $V_H$  or  $V_L$  chains or combinations thereof on the surface of the phage or with aptamers;
- (b) identifying phage or aptamers that bind to said weak D polypeptide of (a); and optionally
  - (c) repeating steps (a) and (b) one or more times.
40. A method of identifying a monoclonal antibody specifically binding to a weak D polypeptide/antigen of claim 13 comprising
- (a) contacting the weak D polypeptide and
  - (aa) a second or more weak D polypeptide(s) and/or
  - (ab) a normal D polypeptide
- wherein the second or more weak D polypeptide(s) and/or the normal D polypeptide are present in a molar mass which is higher, equal or less than the weak D polypeptide of (a) with one or more monoclonal antibodies;
- (b) identifying monoclonal antibodies that bind to said weak D polypeptide of (a); and optionally
  - (c) repeating steps (a) and (b) one or more times.
41. The method according to any one of claims 37 to 40, wherein the weak D polypeptide is exposed on the surface of a cell.

42. The method according to any one of claims 37 to 41, wherein the polypeptide or host cell is affixed to a solid support.
43. The method of any one of claims 37 to 42, wherein subsequent to step (b) or (c), the following step is carried out:
  - (d) identifying the amino acid sequence of the  $V_H$  or  $V_L$  chains and/or identifying the nucleic acid sequence encoding said amino acid sequence.
44. The method according to any one of claims 39 to 43, wherein, in the case that only one round of selection is employed for the identification, the number of weak D polypeptide molecules of (a) is in molar excess over the number of phage particles.
45. Use of cells, preferably red blood cells, from probands for the assessment of the affinity, avidity and/or reactivity of monoclonal anti-D antibodies or of polyclonal anti-D antisera or of anti-globulin or of anti-human-globulin antisera or of preparations thereof.
46. A method for determining whether a patient in need of a blood transfusion is to be transfused with Rh D negative blood from a donor comprising the step of testing a sample from said patient for the presence of one or more Rh D antigens of claim 13, wherein a positive testing for at least one of said antigens is indicative of the need for a transfusion with Rh D negative blood.
47. A method for determining whether blood of a donor may be used for transfusion to a patient in need thereof comprising the step of testing a sample from said donor for the presence of one or more Rh D antigens of claim 13, wherein a positive testing for at least one of said antigens excludes the transfusion of patients that are typed as having wild type Rh D antigen or (a) weak D type(s) other than the weak D type(s) of said donor.

48. Kit comprising
- (a) the oligonucleotide of claim 14; and/or
  - (b) the antibody of claim 15; and/or
  - (c) the antibody of claim 16;
  - (d) the aptamer of claim 15 or 16; and/or
  - (e) the phage of claim 15 or 16.

1/5

Figure 1:





2 / 5

Figure 2.

1	20	40	60
ATG AGC TCT AAG TAC CCG CGG TCT GTC CGG CGC TGC CTG CCC CTC TGG GCC CTA ACA CTG			
Met Ser Ser Lys Tyr Pro Arg Ser Val Arg Arg Cys Leu Pro Leu Trp Ala Leu Thr Leu			
1	10		20
80	100		120
GAA GCA GCT CTC ATT CTC CTC TTC TAT TTT TTT ACC CAC TAT GAC GCT TCC TTA GAG GAT			
Glu Ala Ala Leu Ile Leu Leu Phe Tyr Phe Phe Thr His Tyr Asp Ala Ser Leu Glu Asp			
	30		40
140	160		180
CAA AAG GGG CTC GTG GCA TCC TAT CAA GTT GGC CAA GAT CTG ACC GTG ATG GCG GCC ATT			
Gln Lys Gly Leu Val Ala Ser Tyr Gln Val Gly Gln Asp Leu Thr Val Met Ala Ala Ile			
	50		60
200	220		240
GGC TTG GGC TTC CTC ACC TCG AGT TTC CGG AGA CAC AGC TGG AGC AGT GTG GCC TTC AAC			
Gly Leu Gly Phe Leu Thr Ser Ser Phe Arg Arg His Ser Trp Ser Ser Val Ala Phe Asn			
	70		80
260	280		300
CTC TTC ATG CTG GCG CTT GGT GTG CAG TGG GCA ATC CTG CTG GAC GGC TTC CTG AGC CAG			
Leu Phe Met Leu Ala Leu Gly Val Gln Trp Ala Ile Leu Leu Asp Gly Phe Leu Ser Gln			
	90		100
320	340		360
TTC CCT TCT GGG AAG GTG GTC ATC ACA CTG TTC AGT ATT CGG CTG GCC ACC ATG AGT GCT			
Phe Pro Ser Gly Lys Val Val Ile Thr Leu Phe Ser Ile Arg Leu Ala Thr Met Ser Ala			
	110		120
380	400		420
TTG TCG GTG CTG ATC TCA GTG GAT GCT GTC TTG GGG AAG GTC AAC TTG GCG CAG TTG GTG			
Leu Ser Val Leu Ile Ser Val Asp Ala Val Leu Gly Lys Val Asn Leu Ala Gln Leu Val			
	130		140
440	460		480
GTG ATG GTG CTG GTG GAG GTG ACA GCT TTA GGC AAC CTG AGG ATG GTC ATC AGT AAT ATC			
Val Met Val Leu Val Glu Val Thr Ala Leu Gly Asn Leu Arg Met Val Ile Ser Asn Ile			
	150		160
500	520		540
TTC AAC ACA GAC TAC CAC ATG AAC ATG ATG CAC ATC TAC GTG TTC GCA GCC TAT TTT GGG			
Phe Asn Thr Asp Tyr His Met Asn Met Met His Ile Tyr Val Phe Ala Ala Tyr Phe Gly			
	170		180
560	580		600
CTG TCT GTG GCC TGG TGC CTG CCA AAG CCT CTA CCC GAG GGA ACG GAG GAT AAA GAT CAG			
Leu Ser Val Ala Trp Cys Leu Pro Lys Pro Leu Pro Glu Gly Thr Glu Asp Lys Asp Gln			
	190		200
620	640		660
ACA GCA ACG ATA CCC AGT TTG TCT GCC ATG CTG GGC GCC CTC TTC TTG TGG ATG TTC TGG			
Thr Ala Thr Ile Pro Ser Leu Ser Ala Met Leu Gly Ala Leu Phe Leu Trp Met Phe Trp			
	210		220

3 / 5

Figure 2. Continued.

---

680	700	720
CCA AGT TTC AAC TCT GCT CTG CTG AGA AGT CCA ATC GAA AGG AAG AAT GCC GTG TTC AAC		
Pro Ser Phe Asn Ser Ala Leu Leu Arg Ser Pro Ile Glu Arg Lys Asn Ala Val Phe Asn		
230		240
740	760	780
ACC TAC TAT GCT GTA GCA GTC AGC GTG GTG ACA GCC ATC TCA GGG TCA TCC TTG GCT CAC		
Thr Tyr Tyr Ala Val Ala Val Ser Val Val Thr Ala Ile Ser Gly Ser Ser Leu Ala His		
250		260
800	820	840
CCC CAA GGG AAG ATC AGC AAG ACT TAT GTG CAC AGT GCG GTG TTG GCA GGA GGC GTG GCT		
Pro Gln Gly Lys Ile Ser Lys Thr Tyr Val His Ser Ala Val Leu Ala Gly Gly Val Ala		
270		280
860	880	900
GTG GGT ACC TCG TGT CAC CTG ATC CCT TCT CCG TGG CTT GCC ATG GTG CTG GGT CTT GTG		
Val Gly Thr Ser Cys His Leu Ile Pro Ser Pro Trp Leu Ala Met Val Leu Gly Leu Val		
290		300
920	940	960
GCT GGG CTG ATC TCC GTC GGG GGA GCC AAG TAC CTG CCG GGG TGT TGT AAC CGA GTG CTG		
Ala Gly Leu Ile Ser Val Gly Gly Ala Lys Tyr Leu Pro Gly Cys Cys Asn Arg Val Leu		
310		320
980	1000	1020
GGG ATT CCC CAC AGC TCC ATC ATG GGC TAC AAC TTC AGC TTG CTG GGT CTG CTT GGA GAG		
Gly Ile Pro His Ser Ser Ile Met Gly Tyr Asn Phe Ser Leu Leu Gly Leu Leu Gly Glu		
330		340
1040	1060	1080
ATC ATC TAC ATT GTG CTG CTG GTG CTT GAT ACC GTC GGA GCC GGC AAT GGC ATG ATT GGC		
Ile Ile Tyr Ile Val Leu Leu Val Leu Asp Thr Val Gly Ala Gly Asn Gly Met Ile Gly		
350		360
1100	1120	1140
TTC CAG GTC CTC CTC AGC ATT GGG GAA CTC AGC TTG GCC ATC GTG ATA GCT CTC ACG TCT		
Phe Gln Val Leu Leu Ser Ile Gly Glu Leu Ser Leu Ala Ile Val Ile Ala Leu Thr Ser		
370		380
1160	1180	1200
GGT CTC CTG ACA GGT TTG CTC CTA AAT CTT AAA ATA TGG AAA GCA CCT CAT GAG GCT AAA		
Gly Leu Leu Thr Gly Leu Leu Leu Asn Leu Lys Ile Trp Lys Ala Pro His Glu Ala Lys		
390		400
1220	1240	1251
TAT TTT GAT GAC CAA GTT TTC TGG AAG TTT CCT CAT TTG GCT GTT GGA TTT TAA		
Tyr Phe Asp Asp Gln Val Phe Trp Lys Phe Pro His Leu Ala Val Gly Phe ***		
410		417


---

Figure 3

		10	20	30	40	50	
RHCE	AGCCACTTCA	ACGTTTGTAG	TCTCAGTGGC	CTCATCTGTA	AAGTGAGAAT		650
RHD	-----	-----	-----	-----	-----G-----		
RHCE	TAAGAGATGG	TGCATGTAAA	GTGCTTAACG	GGGAGTAAAT	GGTAGGCAAA		700
RHD	-----	-----	-----	-----	-----		
RHCE	CATTAGCTGC	TGCTATTAGT	ACAGAGAGAC	AATGGTGTGT	GTGAGTCTTG		750
RHD	-----	-----	-A-----	G-----	-----		
RHCE	TGGGCAGAGA	TGGGTGAGAG	GGGAGACAAA	ACAAGTTCTC	ATGATGATGG		800
RHD	-----	-----	-----	-----	-----		
RHCE	GGGCAGGGGG	TCCAGCTGGT	GGTGTCTGGAG	GGAAGTCTGG	ACAGACCCAGT		850
RHD	---A-----C	-----	-----	-----	-----		
		*****	*****	*****	*****		
RHCE	GGTGGGGCTC	GGGTGGGAGG	CACTGGGGGG	GCTGGAGTGG	AAAGAATGTG		900
RHD	-----	-----	-----	-----	-----		
	*****	*****	*****	*****	*****		
RHCE	GCCACAGATG	ACAGCTTCAC	AGCAGAATTC	AGTGCTAAGA	GGAAGTGAGT		950
RHD	-----	-----	-----	-----	-----		
	*****	*****	*****	*****	*****		
RHCE	GGCCATGAGT	TCCATGGTGA	CAGAAAGTCT	AAGACACCTA	GCAAGGCAGG		1000
RHD	-----	-----	-----	-----C-----	-----		
	*****	*****	*****	*****			
RHCE	AGTGGGTGTC	AGCTCAGGGA	AGCTCAGAGG	CTAAACCTAG	GTGAGAGCTG		1050
RHD	-----	-A-----	---C-----	---T-----	-----		
RHCE	AGGGTGTCTAG	ATAAGAGCAA	GGCAAGGCTC	CGGTTCTGGA	GTAAGTGAAGG		1100
RHD	-----	-----	-----	-----	-C-----		
RHCE	ACATAGCAGA	GCTATAACCC	AGGAACAAGG	CCCAGCTTAT	TGGAAGTGGG		1150
RHD	-----	-----G-----	-----	-----	---A-----		
RHCE	ACCAATCACA	CAGGGTGGCA	CAGGCACCAA	GTAAGCAATA	ATAATAATAA		1200
RHD	C-----	-----	-----	-----	-----		
RHCE	AAACAATAAC	AATGATTTAT	GTCTATTGGG	CATTTATTCA	TGTTCTATGC		1250
RHD	-----	-----G-----	---C-----	-----	-----		
RHCE	CAGACACTGG	ACTAAGAGCT	TTATATGTGG	AAACTCATTT	AATCCTTACA		1300
RHD	-----	G-----	-----	-----	-----		

5/5

Figure 4:



# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/08319

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/47 C12Q1/68 C07K16/18 G01N33/566

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N C07K C12Q G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ROUILLAC C ET AL: "Molecular basis of the altered antigenic expression of RhD in weak D (Du) and RhC/e in Rn phenotypes" BLOOD, vol. 87, no. 11, 1 June 1996 (1996-06-01), pages 4853-4861, XP002108679 cited in the application page 4853, last paragraph; table 1 ---	1-3,9, 14-26, 45-48
X	MAAS JH ET AL.: "Rh-D-Genotypisierung auf Exon 2, 5 und 7 von deutschen und japanischen Blutspendern mit sequenzspezifischer Polymerasekettenreaktion" BEITRÄGE ZUR INFUSIONSTHERAPIE UND TRANSFUSIONSMEDIZIN, vol. 34, 1997, pages 203-209, XP002108678 paragraph bridging pages 204 and 205 --- -/-	1-3,9, 14, 17-26,48

☒ Further documents are listed in the continuation of box C.

☐ Patent family members are listed in annex.

\* Special categories of cited documents:

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"Z" document member of the same patent family

Date of the actual completion of the international search

9 July 1999

Date of mailing of the international search report

22/07/1999

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Cupido, M

# INTERNATIONAL SEARCH REPORT

International Application No

PCT/EP 98/08319

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>AVENT ND ET AL: "Evidence of genetic diversity underlying Rh D-, weak D (Du), and partial D phenotypes as determined by multiplex polymerase chain reaction analysis of the RHD gene"  BLOOD,  vol. 89, no. 7, 1 April 1997 (1997-04-01),  pages 2568-2577, XP002108680  cited in the application  table 2</p>	<p>1-3,  9-14,  17-26, 48</p>
P, X	<p>LEGLER TJ ET AL: "RHD genotyping in weak D phenotypes by multiple polymerase chain reactions"  TRANSFUSION,  vol. 38, May 1998 (1998-05), pages  434-440, XP002108681  the whole document</p>	<p>1-48</p>
T	<p>WAGNER FF ET AL: "Molecular basis of weak D phenotypes"  BLOOD,  vol. 93, no. 1,  1 January 1999 (1999-01-01), pages  385-393, XP002108682  the whole document</p>	<p>1-48</p>